

THE CLASSIFYING LIE ALGEBROID OF A GEOMETRIC STRUCTURE I: CLASSES OF COFRAMES

RUI LOJA FERNANDES AND IVAN STRUCHINER

CONTENTS

1. Introduction	1
2. Equivalence of coframes and realization problems	6
2.1. Local equivalence	6
2.2. Global equivalence	8
3. The classifying algebroid	10
3.1. Lie algebroids and Cartan's Problem	10
3.2. Realizations and Lie algebroid morphisms	12
4. Maurer-Cartan Forms on Lie groupoids	13
4.1. Definition of a Maurer-Cartan Form	14
4.2. The Local Universal Property	15
4.3. The Global Universal Property	17
5. Local Classification and symmetries	19
5.1. Local Classification Problem	19
5.2. Equivalence versus formal equivalence	20
5.3. Symmetries of Realizations	21
6. Global Equivalence	24
6.1. The Globalization Problem	24
6.2. The Global Classification Problem	26
6.3. Cohomological Invariants of Geometric Structures	28
6.4. The modular class	30
7. An Example: Surfaces of Revolution	32
Appendix A. The classifying Lie algebroid of a single coframe	36
References	39

1. INTRODUCTION

This is the first of two papers dedicated to a systematic study of symmetries, invariants and moduli spaces of geometric structures of *finite type*. Roughly speaking, a geometric structure of finite type can be described as any geometric structure on a smooth manifold which determines (and is determined by) a coframe on a (possibly different) manifold. Here by a **coframe** on an n -dimensional manifold M we mean a generating set $\{\theta^1, \dots, \theta^n\}$ of everywhere linearly independent 1-forms on M .

Examples of finite type geometric structures include finite type G -structures (see, e.g., [19] or [10]), Cartan geometries (see, e.g., [17] or [10]) or linear connections which preserve some tensor on a manifold (see, e.g., [11]).

The main problem concerning finite type structures is that of determining when two such geometric structures are isomorphic. This is a classical problem, that goes back to Élie Cartan [3], and is usually refer to as the *Cartan equivalence problem*. When two geometric structures of the same kind have been properly characterized

by coframes $\{\theta^i\}$ and $\{\bar{\theta}^i\}$ on n -dimensional manifolds M and \bar{M} , the equivalence problem takes the form:

Problem 1.1 (Equivalence Problem). *Does there exist a diffeomorphism $\phi : M \rightarrow \bar{M}$ satisfying*

$$\phi^* \bar{\theta}^i = \theta^i$$

for all $1 \leq i \leq n$?

We will show that to a coframe there is associated a **classifying Lie algebroid** which encodes the corresponding equivalence problem. This classifying algebroid plays a crucial role in various geometric, local and global, classification problems associated with coframes.

In order to explain how this algebroid arises, recall that the solution to the equivalence problem is based on the simple fact that exterior differentiation and pullbacks commute: taking exterior derivatives and using the fact that $\{\theta^i\}$ is a coframe, we may write the **structure equations**

$$(1.1) \quad d\theta^k = \sum_{i < j} C_{ij}^k(x) \theta^i \wedge \theta^j$$

for uniquely defined functions $C_{ij}^k \in C^\infty(M)$ known as **structure functions**. Analogously, we may write

$$d\bar{\theta}^k = \sum_{i < j} \bar{C}_{ij}^k(\bar{x}) \bar{\theta}^i \wedge \bar{\theta}^j.$$

It then follows from $\phi^* d\bar{\theta}^k = d\phi^* \bar{\theta}^k$, that a necessary condition for equivalence is that

$$\bar{C}_{ij}^k(\phi(x)) = C_{ij}^k(x).$$

Thus, the structure functions are *invariants* of the coframe. One can obtain more invariants of the coframe by further differentiation and this is the key remark to solve the equivalence problem.

In the opposite direction, one may consider the problem of determining when a given set of functions C_{ij}^k can be realized as the structure functions of some coframe. This problem was proposed and solved by Cartan in [3]. Following Bryant (see the Appendix in [1]), we state it in the following precise form:

Problem 1.2 (Cartan's Realization Problem). *One is given:*

- an integer $n \in \mathbb{N}$,
- a d -dimensional manifold X ,
- a set of functions $C_{ij}^k \in C^\infty(X)$ with indexes $1 \leq i, j, k \leq n$,
- and n vector fields $F_i \in \mathfrak{X}(X)$

and asks for the existence of

- an n -dimensional manifold M ;
- a coframe $\theta = \{\theta^i\}$ on M ;
- a map $h : M \rightarrow X$

satisfying

$$(1.2) \quad d\theta^k = \sum_{i < j} C_{ij}^k(h(m)) \theta^i \wedge \theta^j$$

$$(1.3) \quad dh = \sum_i F_i(h(m)) \theta^i.$$

Note that, most often, in concrete examples one is given the structure functions and looks for all the possible coframes realizing it. In fact, one is interested in the following two problems:

Local Classification Problem: What are all germs of coframes which solve Cartan's realization problem?

Local Equivalence Problem: When are two such germs of coframes equivalent?

Later, we shall give complete solutions to the existence problem, as well as both the classification and the equivalence problems. Before we proceed, let us present a simple, classic, but elucidating example showing that Lie theory enters into the picture.

Example 1.3 (The case $d = 0$). *Suppose that we are interested in the equivalence and classification problem for coframes whose structure functions are constant, or in the precise formulation of Cartan's realization problem given above, the case where the manifold X reduces to a point (in particular, the vector fields F_i vanish identically).*

Necessary conditions for the existence of a solution to the realization problem are obtained from

$$d^2\theta^m = 0 \quad (m = 1, \dots, n).$$

These conditions imply that the the structure constants must satisfy:

$$C_{im}^l C_{jk}^m + C_{jm}^l C_{ki}^m + C_{km}^l C_{ij}^m = 0, \quad (i, j, k, m = 1, \dots, n),$$

i.e., they are the structure constants of a Lie algebra \mathfrak{g} relative to same basis $\{e_1, \dots, e_n\}$:

$$[e_i, e_j] = C_{ij}^k e_k.$$

This condition is also sufficient. In fact, if we let $M := G$ be any Lie group which has \mathfrak{g} as its Lie algebra, then the components θ^i of its \mathfrak{g} -valued right invariant Maurer-Cartan form ω_{MC} relative to some basis $\{e_1, \dots, e_n\}$ of \mathfrak{g} , have C_{ij}^k as its structure functions.

Now let (M, θ^i, h) be any other solution of the realization problem. If we define a \mathfrak{g} -valued 1-form on M by

$$\theta = \sum_i \theta^i e_i,$$

then the structure equation (1.2) for the coframe $\{\theta^i\}$ implies that θ satisfies the Maurer-Cartan equation

$$d\theta + \frac{1}{2}[\theta, \theta] = 0.$$

It is then well known that there exists a locally defined diffeomorphism $\psi : M \rightarrow G$ onto a neighborhood of the identity such that

$$\theta = \psi^* \omega_{MC}.$$

This is sometimes referred to as the universal property of the Maurer-Cartan form. Thus, at least locally, there is only one solution to the realization problem, up to equivalence.

In general, when the structure functions are not constant, they cannot determine a Lie algebra. Instead, we have the following result which is the basic proposition which underlies all our approach:

Theorem 1.4. *Solutions to Cartan's problem exist if and only if the initial data to the problem determines a Lie algebroid structure on the trivial vector bundle $A \rightarrow X$ with fiber \mathbb{R}^n , Lie bracket $[\cdot, \cdot]_A : \Gamma(A) \times \Gamma(A) \rightarrow \Gamma(A)$ relative to the standard basis of sections $\{e_1, \dots, e_n\}$ given by:*

$$[e_i, e_j]_A = C_{ij}^k(x) e_k,$$

and anchor $\sharp : A \rightarrow TM$ defined by $\sharp(e_i) = F_i$.

We will call $A \rightarrow X$ the **classifying Lie algebroid** of the Cartan's realization problem. Also, the map $h : M \rightarrow X$, which determines the particular coframe we are interested in, will be called the **classifying map** of the realization.

We will see in Section 3.1, that given a coframe θ on a manifold M , there is associated to it a Cartan Realization Problem, and hence a classifying Lie algebroid A_θ . This will be called the **classifying Lie algebroid of the coframe**. It is important to make the following distinctions:

- (i) To a *single* coframe θ , there is associated a Cartan realization problem and, hence, a classifying Lie algebroid A_θ .
- (ii) To a Cartan realization problem, there is associated a classifying Lie algebroid A and a *family* of coframes (its solutions).

Note that distinct Cartan realization problems can share the same coframe as solutions. However, we have the following:

Proposition 1.5. *Let (n, X, C_{ij}^k, F_i) be a Cartan realization problem with associated classifying algebroid A and let (M, θ, h) be a connected realization. Then $h(M)$ is an open subset of an orbit of A so that $A|_{h(M)} \rightarrow h(M)$ is also a Lie algebroid. Moreover, there exists a Lie algebroid morphism from $A|_{h(M)}$ to A_θ (the classifying algebroid of the coframe θ) which is a fiberwise isomorphism that covers a submersion $h(M) \rightarrow X_\theta$.*

In general, a Lie algebroid does not need to be associated with any Lie groupoid (see [5] for a complete discussion of this point). However, assume for now that this is the case for the classifying Lie algebroid $A \rightarrow X$ and denote by $\mathcal{G} \rightrightarrows X$ a Lie groupoid integrating it. One can then introduce the Maurer-Cartan form ω_{MC} on \mathcal{G} and it follows from its universal property that:

Theorem 1.6. *Let ω_{MC} be the Maurer-Cartan form on a Lie groupoid \mathcal{G} integrating the classifying Lie algebroid A of a Cartan's realization problem. Then the pull-back of ω_{MC} to each \mathfrak{s} -fiber induces a coframe which gives a solution to the realization problem. Moreover, every solution to Cartan's problem is locally equivalent to one of these.*

This settles the classification and the equivalence problems. It also shows one of the main advantages of using our Lie algebroid approach: the classifying Lie algebroid (or more precisely its local Lie groupoid) gives rise to a recipe for constructing explicit solutions to the problem. The solutions are the \mathfrak{s} -fibers of the groupoid equipped with the restriction of the Maurer-Cartan forms. There are several other advantages in using this Lie algebroid approach.

First of all, it turns out that properties of the classifying Lie algebroid are closely related to geometric properties of the coframes it classifies. For example, let us call any self-equivalence of a coframe which preserves the classifying map h a **symmetry of the realization**. Then:

Proposition 1.7. *Let $A \rightarrow X$ be the classifying Lie algebroid of a Cartan's realization problem. Then:*

- (1) *To every point on X there corresponds a germ of a coframe which is a solution to the realization problem;*
- (2) *Two such germs of realizations are equivalent if and only if they correspond to the same point in X ;*
- (3) *The isotropy Lie algebra of $A \rightarrow X$ at some $x \in X$ is isomorphic to the symmetry Lie algebra of the corresponding germ of realization.*

Second of all, the Lie algebroid approach is well fitted to the study of global aspects of the theory. The first of these global issues that we will consider is the

Problem 1.8 (Globalization Problem). *Given a Cartan's problem with initial data (n, X, C_{ij}^k, F_i) and two germs of coframes θ_0 and θ_1 which solve the problem, does there exist a global connected solution (M, θ, h) to the realization problem for which θ_0 is the germ of θ at a point $p_0 \in M$ and θ_1 is the germ of θ at a point $p_1 \in M$?*

Another important global issue is the *global equivalence problem*. In general, the classifying Lie algebroid does not distinguish between a realization and its covers. Here, by a **realization cover** of (M, θ, h) we mean a realization of the form $(\tilde{M}, \pi^*\theta, \pi^*h)$ where $\pi : \tilde{M} \rightarrow M$ is a covering map.

It is then natural to consider the following equivalence relation on the set of realizations of a Cartan's problem. Two realization (M_1, θ_1, h_1) and (M_2, θ_2, h_2) are said to be **globally equivalent, up to covering** if they have a common realization cover (M, θ, h) , i.e.,

$$\begin{array}{ccc} & (M, \theta, h) & \\ \pi_1 \swarrow & & \searrow \pi_2 \\ (M_1, \theta_1, h_1) & & (M_2, \theta_2, h_2). \end{array}$$

This leads to:

Problem 1.9 (Global Classification Problem). *What are all the solutions of a Cartan's realization problem up to global equivalence, up to covering?*

Of course, the solutions to these global problems rely, again, on understanding the classifying Lie algebroid. We need to assume that the restriction of the classifying Lie algebroid to each orbit comes from a Lie groupoid. Note that this assumption is weaker than requiring integrability of the whole Lie algebroid A (see [5]), so in this case we will say that A is *weakly integrable*.

For the globalization problem we then have the following solution:

Theorem 1.10. *Let $A \rightarrow X$ be the classifying Lie algebroid of a Cartan's realization problem. Two germs of coframes which belong to the same global connected realization correspond to points in the same orbit of A . Moreover, if A is weakly integrable then the converse is also true: two germs of coframes which correspond to points in the same orbit belong to the same global connected realization.*

On the other hand, for the global equivalence problem we obtain the following result which generalizes the two main theorems concerning global equivalences due to Olver [13]:

Theorem 1.11. *Assume that the classifying Lie algebroid $A \rightarrow X$ of a Cartan's realization problem is weakly integrable. Then any solution to Cartan's problem is globally equivalent, up to covering, to an open set of an \mathfrak{s} -fiber of the source 1-connected Lie groupoid integrating $A|_L$, for some leaf of A .*

Finally, the classifying Lie algebroid can also be used to produce invariants of a geometric structure of finite type, as well as to recover classical results about their symmetry groups. In fact, if the geometric structure is characterized by a coframe θ on a manifold M , with associated classifying algebroid A_θ , then we can view the coframe as a Lie algebroid morphism $\theta : TM \rightarrow A_\theta$ (which covers the classifying map h). As a general principle, the coframe (viewed as a Lie algebroid morphism) pulls back invariants of the classifying Lie algebroid A_θ to invariants of the coframe. To illustrate this point of view we will introduce a new invariant called the *modular class of a coframe* and which is obtained as the pullback of the modular class of its classifying Lie algebroid.

We can summarize this outline of the paper by saying that there are essentially two ways in which we use the existence of a classifying Lie algebroid for a Cartan's realization problem.

- For a *class* of geometric structures of finite type whose moduli space (of germs) is finite dimensional, we can set up a Cartan's realization problem whose classifying Lie algebroid gives us information about its local equivalence and classification problems, as well as its globalization and global classification problems.
- For a *single* geometric structure of finite type, we can set up a Cartan's realization problem whose classifying Lie algebroid describes the symmetries of the geometry, and also provides a recipe for producing invariants of the structure.

In a sequel to this paper [10], we will discuss finite type G -structures and show how one can associate a classifying Lie algebroid to such a structure. There we will also present several geometric examples related to torsion-free connections on G -structures.

Part of the results described in this paper were obtained by the second author in his PhD Thesis [20].

2. EQUIVALENCE OF COFRAMES AND REALIZATION PROBLEMS

In this section, we will explain how coframes give rise to Cartan realization problems. First we consider the local problem, where we assume the coframe to be fully regular at some point. Then we consider the global problem for coframes which are fully regular at every point. Our exposition maybe seen as a Lie algebroid version of the classical approach, which can be found in the monographs of Olver [13] and Sternberg [19], to which we refer the reader for more details.

2.1. Local equivalence. Let $\{\theta^i\}$ and $\{\bar{\theta}^i\}$ be coframes on two n -dimensional manifolds M and \bar{M} . Throughout this text we will use unbarred letters to denote objects on M and barred letters to denote objects on \bar{M} . Any (locally defined) diffeomorphism $\phi : M \rightarrow \bar{M}$ such that $\phi^*\bar{\theta}^i = \theta^i$ will be called an **(local) equivalence**.

As we saw in the Introduction, the structure equations (1.1)

$$d\theta^k = \sum_{i < j} C_{ij}^k(p)\theta^i \wedge \theta^j$$

and the structure functions $C_{ij}^k \in C^\infty(M)$ play a crucial role in the study of the equivalence problem. For example, the structure functions of any coframe $\bar{\theta}$ equivalent to θ must satisfy

$$\bar{C}_{ij}^k(\phi(p)) = C_{ij}^k(p),$$

so we may use the structure functions to obtain a set of necessary conditions for solving the equivalence problem. They are examples of *invariant functions*:

Definition 2.1. A function $I \in C^\infty(M)$ is called an **invariant function of a coframe** $\theta = \{\theta^i\}$ if for any locally defined self equivalence $\phi : U \rightarrow V$, where $U, V \subset M$ are open sets, one has

$$I \circ \phi = I.$$

We denote by $\text{Inv}(\theta) \subset C^\infty(M)$ the space of invariant functions of the coframe θ .

Now, for any function $f \in C^\infty(M)$ and coframe θ one can define the coframe derivatives $\frac{\partial f}{\partial \theta^k} \in C^\infty(M)$ as being the coefficients of the differential of f when expressed in terms of the coframe $\{\theta^i\}$,

$$df = \sum_k \frac{\partial f}{\partial \theta^k} \theta^k.$$

Using the fact that $d\phi^* = \phi^*d$, it follows that:

Lemma 2.2. *If $I \in \text{Inv}(\theta)$ is an invariant function of a coframe $\theta = \{\theta^i\}$, then so is $\frac{\partial I}{\partial \theta^k}$ for all $1 \leq k \leq n$.*

Obviously, if $\phi : M \rightarrow \bar{M}$ is an equivalence between coframes $\theta = \{\theta^i\}$ and $\bar{\theta} = \{\bar{\theta}^i\}$, we obtain a bijection $\phi^* : \text{Inv}(\bar{\theta}) \rightarrow \text{Inv}(\theta)$. So, for example, we have as necessary conditions for equivalence that the structure functions and its coframe derivatives of any order

$$C_{ij}^k, \frac{\partial C_{ij}^k}{\partial \theta^l}, \dots, \frac{\partial^s C_{ij}^k}{\partial \theta^{l_1} \dots \partial \theta^{l_s}}, \dots$$

must correspond under the equivalence. This gives an infinite set of conditions for equivalence. However, observe that if invariants $f_1, \dots, f_l \in \text{Inv}(\theta)$ satisfy a functional relationship $f_l = H(f_1, \dots, f_{l-1})$, then the corresponding elements $\bar{f}_1, \dots, \bar{f}_l$ in $\text{Inv}(\bar{\theta})$ must satisfy the same functional relation

$$\bar{f}_l = H(\bar{f}_1, \dots, \bar{f}_{l-1})$$

for the same function H . This shows that we do not need to deal with all the invariant functions in $\text{Inv}(\theta)$, but only with those that are functionally independent.

For any subset $\mathcal{C} \subset C^\infty(M)$ we define the **rank of \mathcal{C} at $p \in M$** , denoted by $r_p(\mathcal{C})$, to be the dimension of the vector space spanned by $\{d_p f : f \in \mathcal{C}\}$. Also, we say that \mathcal{C} is **regular at p** if $r_{p'}(\mathcal{C}) = r_p(\mathcal{C})$ for all p' in a neighborhood of p in M . Finally, we say that \mathcal{C} is **regular**, if it is regular at every point $p \in M$.

Definition 2.3. *A coframe θ is called **fully regular at $p \in M$** if the set $\text{Inv}(\theta)$ is regular at p . It is called **fully regular** if it is fully regular at every $p \in M$.*

By the implicit function theorem, if θ is a fully regular coframe in M of rank d at $p \in M$, then we can find invariant functions $\{h_1, \dots, h_d\} \subset \text{Inv}(\theta)$, independent in a neighborhood of p , such that every $f \in \text{Inv}(\theta)$ can be written as

$$f = H(h_1, \dots, h_d)$$

in a neighborhood of p . In particular, we can assume that there is an open set $U \subset M$ containing p , where this independent set of invariant functions determine a smooth map:

$$h : U \subset M \rightarrow \mathbb{R}^d, \quad h(p) = (h_1(p), \dots, h_d(p)),$$

onto an open set $X \subset \mathbb{R}^d$. Note that $C_{ij}^k = C_{ij}^k(h_1, \dots, h_d)$, so the structure functions are smooth functions $C_{ij}^k \in C^\infty(X)$. Also, differentiating h_a , we find

$$dh_a = \sum_i (F_i^a \circ h) \theta^i,$$

for some smooth functions $F_i^a \in C^\infty(X)$, so we obtain n vector fields on X :

$$F_i = \sum_a F_i^a \frac{\partial}{\partial h_a}.$$

Since the functions C_{ij}^k and the vector fields F_i satisfy

$$\begin{aligned} d\theta^k &= \sum_{i < j} C_{ij}^k(h)\theta^i \wedge \theta^j \\ dh &= \sum_i F_i(h)\theta^i \end{aligned}$$

we conclude that a coframe which is fully regular at p determines the initial data of a Cartan's Realization Problem (Problem 1.2).

Remark 2.4. *The data depends on the choice of the functions $\{h_1, \dots, h_d\} \subset \text{Inv}(\theta)$. However, if $\{h'_1, \dots, h'_d\} \subset \text{Inv}(\theta)$ is a different choice, defining an open set X' , giving rise to data $C'_{ij}^k \in C^\infty(X')$ and $F'_i \in \mathfrak{X}(X')$, then there is a diffeomorphism $\phi : X \rightarrow X'$ defined from a neighborhood of $h(p)$ to a neighborhood of $h'(p)$, which makes the diagram commutative:*

$$\begin{array}{ccc} & U & \\ h \swarrow & & \searrow h' \\ X & \xrightarrow{\phi} & X' \end{array}$$

and is such that $\phi^* C'_{ij}^k = C_{ij}^k$ and $\phi_* F_i = F'_i$.

2.2. Global equivalence. So far, we have only considered a neighborhood of a point where the coframe is fully regular. We consider now the case where θ is a fully regular coframe at every point of M .

Definition 2.5. *We will say that two points p and q of M are **locally formally equivalent** if there exist neighborhoods U_p and U_q of p and q in M and a diffeomorphism $\phi : U_p \rightarrow U_q$ such that $\phi(p) = q$ and*

$$(2.1) \quad f(\phi(p')) = f(p'), \quad \text{for all } p' \in U_p \text{ and } f \in \text{Inv}(\theta).$$

Remark 2.6. *We note that p and q are locally formally equivalent if and only if $f(p) = f(q)$ for all $f \in \text{Inv}(\theta)$. The crucial point is that since $\text{Inv}(\theta)$ is closed under the operation of taking coframe derivatives, it follows that if $f(p) = f(q)$ for all $f \in \text{Inv}(\theta)$, and $f_1, \dots, f_d \in \text{Inv}(\theta)$ are functionally independent in a neighborhood of p , then they are also functionally independent in some neighborhood of q . We shall see later, in Proposition 5.4, that local formal equivalence and local equivalence actually coincide.*

With this definition, we have:

Proposition 2.7. *Let θ be a fully regular coframe on M of rank d and denote by \sim the local formal equivalence relation. The quotient:*

$$X_\theta := M / \sim$$

has a natural structure of a smooth manifold of dimension d .

Proof. On the the orbit space $X_\theta := M / \sim$ we consider the quotient topology and we denote by $\kappa_\theta : M \rightarrow X_\theta$ the projection. Also, if U is an open subset of M , then we will denote by $\text{Inv}_U(\theta)$ the restriction of $\text{Inv}(\theta)$ to U .

We start by choosing a cover of M by a family of open sets $\{U_\lambda\}_{\lambda \in \Lambda}$ such that for each $\lambda \in \Lambda$ there exist elements $h_1^\lambda, \dots, h_d^\lambda \in \text{Inv}(\theta)$ whose restriction to U_λ are functionally independent and generate $\text{Inv}_{U_\lambda}(\theta)$. Thus, on each U_λ we obtain a smooth *open* map $h^\lambda : U_\lambda \rightarrow \mathbb{R}^d$ defined by

$$p \mapsto (h_1^\lambda(p), \dots, h_d^\lambda(p)).$$

The image of this map is an open subset $h^\lambda(U_\lambda) \subset \mathbb{R}^d$, and we have an induced map on the orbit space $\bar{h}^\lambda : X_\lambda \rightarrow \mathbb{R}^d$, where $X_\lambda := \kappa_\theta(U_\lambda) \subset X_\theta$.

Lemma 2.8. *The sets X_λ form an open cover of X_θ and each $\bar{h}^\lambda : X_\lambda \rightarrow \mathbb{R}^d$ is a homeomorphism onto the open set $h^\lambda(U_\lambda) \subset \mathbb{R}^d$.*

Proof of Lemma 2.8. The set $X_\lambda = \kappa_\theta(U_\lambda)$ is open if and only if the saturation of U_λ , defined by

$$\tilde{U}_\lambda := \{q \in M : q \sim p \text{ for some } p \in U_\lambda\},$$

is open in M . This follows because if $q \in \tilde{U}_\lambda$ then there exists $p \in U_\lambda$ and a diffeomorphism $\phi : U_p \rightarrow U_q$, defined on neighborhoods of p and q , such that (2.1) holds. Then $\phi(U_p \cap U_\lambda)$ is an open set containing q , which is contained in \tilde{U}_λ . We conclude that the saturation \tilde{U}_λ is open, therefore X_λ is also open.

The induced map $\bar{h}^\lambda : X_\lambda \rightarrow \mathbb{R}^d$ is continuous (by definition of the quotient topology), open (since h^λ is open) and gives a 1:1 map onto the open set $h^\lambda(U_\lambda) \subset \mathbb{R}^d$. Hence, $\bar{h}^\lambda : X_\lambda \rightarrow h^\lambda(U_\lambda)$ is a homeomorphism. \square

Now observe that if there exists a diffeomorphism defined on an open subset $W_\lambda \subset U_\lambda$ with values in another open subset $W_\mu \subset U_\mu$,

$$\phi : W_\lambda \rightarrow W_\mu,$$

and such that

$$f(\phi(p)) = f(p), \quad \text{for all } p \in W_\lambda, \text{ and } f \in \text{Inv}(\theta),$$

(for example if $U_\lambda \cap U_\mu \neq \emptyset$, and $W_\lambda = U_\lambda \cap U_\mu = W_\mu$) then the restriction of each h_i^μ to W_μ can be written as a smooth function of the (restriction of the) functions $h_1^\lambda, \dots, h_d^\lambda$, i.e.,

$$h_i^\mu = H_{\lambda\mu}^i(h_1^\lambda, \dots, h_d^\lambda).$$

Since both sets $\{h_i^\lambda\}$ and $\{h_i^\mu\}$ are functionally independent, we obtain a diffeomorphism

$$H_{\lambda\mu} : h^\lambda(W_\lambda) \rightarrow h^\mu(W_\mu).$$

Therefore, at the level of the orbit space, we obtain the commutative diagram:

$$\begin{array}{ccc} & X_\lambda \cap X_\mu & \\ \bar{h}^\lambda \swarrow & & \searrow \bar{h}^\mu \\ \bar{h}^\lambda(X_\lambda \cap X_\mu) & \xrightarrow{H_{\lambda\mu}} & \bar{h}^\mu(X_\lambda \cap X_\mu) \end{array}$$

Therefore, the family $\{(X_\lambda, \bar{h}^\lambda)\}$ gives an atlas for X_θ , whose transition functions are precisely the maps $H_{\lambda\mu}$, so we conclude that X_θ is a smooth manifold, possibly non-Hausdorff.

We claim that X_θ is Hausdorff. Let $x, y \in X_\theta$ and suppose that any two neighborhoods of x and y intersect. To prove the claim we must show that $x = y$. For this, pick $p, q \in M$ such that $\kappa_\theta(p) = x$ and $\kappa_\theta(q) = y$. Then, by our assumption, any neighborhoods of p and q contain points which are in the same equivalence class. Therefore, we can choose sequences $p_n \rightarrow p$ and $q_n \rightarrow q$ such that $p_n \sim q_n$. It follows that for any $f \in \text{Inv}(\theta)$:

$$f(p) = \lim f(p_n) = \lim f(q_n) = f(q).$$

This implies that $p \sim q$ (see Remark 2.6) so $x = y$. \square

We will see that the manifold X_θ plays a crucial role in the various equivalence and classification problems associated with a fully regular coframe θ . Hence we suggest the following

Definition 2.9. *The manifold X_θ is called the **classifying manifold** of the coframe and the natural map $\kappa_\theta : M \rightarrow X_\theta$ is called the **classifying map** of the coframe.*

Remark 2.10. *For a arbitrary coframe θ one can still define the (singular) classifying space $X_\theta := M / \sim$, where \sim denotes local equivalence (which, now, may differ from formal local equivalence). The space X_θ is smooth on the dense open set consisting of points where the coframe is fully regular. It is an interesting problem to study the kind of singularities that may arise in this space.*

If $\theta = \{\theta^i\}$ is a fully regular coframe in M of rank d , with classifying manifold X_θ , by Remark 2.4, the structure functions C_{ij}^k can be seen as smooth functions in X_θ , while the locally defined vector fields $F_i = \sum_a F_i^a \frac{\partial}{\partial h_a^x}$ glue to globally defined vector fields $F_i \in \mathfrak{X}(X_\theta)$. We conclude that:

Proposition 2.11. *If θ is a fully regular coframe on a manifold M , then it gives rise to a Cartan's realization problem with initial data $(n, X_\theta, C_{ij}^k, F_i)$ for which $(M, \theta, \kappa_\theta)$ is a solution.*

3. THE CLASSIFYING ALGEBROID

In the previous section we have seen that the equivalence problem for coframes leads naturally to a Cartan realization problem. We now show that if such a problem admits solutions, then it is associated with a Lie algebroid, called the *classifying algebroid* of the realization problem.

3.1. Lie algebroids and Cartan's Problem. Recall that a Lie algebroid $A \rightarrow X$ is a vector bundle equipped with a Lie bracket on its space of sections $\Gamma(A)$ and a bundle map $\sharp : A \rightarrow TX$, called the anchor, such that the following Leibniz type identity holds:

$$[\alpha, f\beta] = f[\alpha, \beta] + (\mathcal{L}_{\sharp(\alpha)}f)\beta, \quad (\alpha, \beta \in \Gamma(A), f \in C^\infty(X)).$$

We refer to [2, 7] for details on Lie algebroids. If the vector bundle A is trivial, so that we have a basis of global sections $\{\alpha_1, \dots, \alpha_n\}$, then we can express the bracket by:

$$(3.1) \quad [\alpha_i, \alpha_j] = \sum_{k=1}^n C_{ij}^k \alpha_k,$$

for some structure functions $C_{ij}^k \in C^\infty(U)$. The anchor, in turn, is characterized by the vector fields $F_i \in \mathfrak{X}(X)$ given by:

$$(3.2) \quad F_i := \sharp(\alpha_i), \quad (i = 1, \dots, n).$$

The Leibniz identity and the Jacobi identity for the bracket, lead to the following set of equations:

$$(3.3) \quad [F_i, F_j] = \sum_k C_{ij}^k F_k$$

$$(3.4) \quad \mathcal{L}_{F_j} C_{kl}^i + \mathcal{L}_{F_k} C_{lj}^i + \mathcal{L}_{F_l} C_{jk}^i = \sum_m (C_{mj}^i C_{kl}^m + C_{mk}^i C_{lj}^m + C_{ml}^i C_{jk}^m)$$

Conversely, if one is given functions $C_{ij}^k \in C^\infty(X)$ and vector fields $F_i \in \mathfrak{X}(X)$ satisfying these equations, then one obtains a Lie algebroid structure on the trivial vector bundle $A := X \times \mathbb{R}^n$ with Lie bracket (3.1) and anchor (3.2).

Remark 3.1. *If one also chooses local coordinates (x_1, \dots, x_d) on a domain $U \subset X$, so that the vector fields F_i are expressed as:*

$$F_i = \sum_a F_i^a \frac{\partial}{\partial x_a},$$

then we obtain the complete set of structure functions $C_{ij}^k, F_i^a \in C^\infty(U)$ of the Lie algebroid A . Equations (3.3) and (3.4) became:

$$\sum_{b=1}^d \left(F_i^b \frac{\partial F_j^a}{\partial x_b} - F_j^b \frac{\partial F_i^a}{\partial x_b} \right) = \sum_{l=1}^n C_{ij}^l F_l^a$$

$$\sum_{b=1}^d \left(F_j^b \frac{\partial C_{kl}^i}{\partial x_b} + F_k^b \frac{\partial C_{lj}^i}{\partial x_b} + F_l^b \frac{\partial C_{jk}^i}{\partial x_b} \right) = \sum_{m=1}^n (C_{mj}^i C_{kl}^m + C_{mk}^i C_{lj}^m + C_{ml}^i C_{jk}^m)$$

*We call this set of equations, or its shorter form (3.3) and (3.4), the **structure equations** of the Lie algebroid.*

We can now give the necessary conditions for solving the realization problem. They are simply a consequence of the fact that $d^2 = 0$:

Proposition 3.2. *Let (n, X, C_{ij}^k, F_i) be the initial data of a Cartan's realization problem. If for every $x_0 \in X$ there is a realization (M, θ, h) with $h(m_0) = x_0$, for some $m_0 \in M$, then the $-C_{ij}^k \in C^\infty(X)$ and $F_i \in \mathfrak{X}(X)$ determine the structure of a Lie algebroid A over X .*

Proof. Differentiating equations (1.2) and (1.3) and using $d^2 = 0$, we obtain:

$$\mathcal{L}_{F_j} C_{kl}^i + \mathcal{L}_{F_k} C_{lj}^i + \mathcal{L}_{F_l} C_{jk}^i = - \sum_m (C_{mj}^i C_{kl}^m + C_{mk}^i C_{lj}^m + C_{ml}^i C_{jk}^m)$$

$$[F_i, F_j] = - \sum_k C_{ij}^k F_k.$$

These are just the structure equations (3.3) and (3.4) for the Lie algebroid defined by the trivial vector bundle $A \rightarrow X$ with fiber \mathbb{R}^n , with Lie bracket and anchor given by:

$$\sharp(\alpha_i) := F_i,$$

$$[\alpha_i, \alpha_j] := - \sum_k C_{ij}^k \alpha_k,$$

where $\{\alpha_1, \dots, \alpha_n\}$ is the canonical basis of sections of A . □

Using this proposition we introduce the following basic concept:

Definition 3.3. *The Lie algebroid constructed out of the initial data of a Cartan's realization problem will be called the **classifying algebroid** of the problem.*

We will see later that the fact that the initial data of a Cartan realization problem determines a Lie algebroid is also sufficient for the existence of solutions, and that this Lie algebroid leads to solutions of the various classifications problems associated to coframes, which justifies its name.

Remark 3.4. *Notice that if we start with a fixed coframe θ on a manifold M , then we have an associated Cartan realization problem. The corresponding Lie algebroid will be called the **classifying algebroid of the coframe** and denoted $A_\theta \rightarrow X_\theta$. If the coframe is also the solution of another realization problem, with associated algebroid A , we will see later (Corollary 5.11) the precise relationship between A_θ and A .*

3.2. Realizations and Lie algebroid morphisms. Let $A \rightarrow X$ be the classifying Lie algebroid of a Cartan's problem. Each realization (M, θ, h) of the problem determines a bundle map:

$$(3.5) \quad \begin{array}{ccc} TM & \xrightarrow{\theta} & A \\ \downarrow & & \downarrow \\ M & \xrightarrow{h} & X \end{array}$$

by setting:

$$\begin{aligned} TM &\longrightarrow A = X \times \mathbb{R}^n, \\ v &\longmapsto (h(p(v)), (\theta^1(v), \dots, \theta^n(v))), \end{aligned}$$

where $p : TM \rightarrow M$ denotes the projection. This bundle map is a fiberwise isomorphism.

We have the following basic result:

Proposition 3.5. *Let $A \rightarrow X$ be the classifying Lie algebroid of a Cartan's problem. The realizations of this problem are in 1:1 correspondence with bundle maps (3.5) which are Lie algebroid morphisms and fiberwise isomorphisms.*

Before we prove this result, let us recall some basic facts about Lie algebroid morphisms (see, e.g., [7]). A Lie algebroid morphism is a bundle map:

$$\begin{array}{ccc} A & \xrightarrow{\mathcal{H}} & B \\ \downarrow & & \downarrow \\ X & \xrightarrow{h} & Y \end{array}$$

which commutes with anchors:

$$dh \circ \sharp_A = \sharp_B \circ \mathcal{H},$$

and which preserves brackets. Since we are dealing with a bundle map over different basis, to express this condition is somewhat cumbersome (see [7]). For us, it will be convenient to use the generalized Maurer-Cartan equation, which we now recall.

First we introduce an arbitrary connection ∇ on the vector bundle $B \rightarrow Y$. Then for *any* bundle maps $\mathcal{H}, \mathcal{G} : A \rightarrow B$ which commute with the anchors, we define:

- (a) The differential operator $d_\nabla \mathcal{H}$: for each $\alpha_1, \alpha_2 \in \Gamma(A)$, $d_\nabla \mathcal{H}(\alpha_1, \alpha_2)$ is the section of the pull-back bundle $\Gamma(h^*B)$ given by¹:

$$(3.6) \quad d_\nabla \mathcal{H}(\alpha_1, \alpha_2) := \nabla_{\sharp_A \alpha_1} \mathcal{H}(\alpha_2) - \nabla_{\sharp_A \alpha_2} \mathcal{H}(\alpha_1) - \mathcal{H}([\alpha_1, \alpha_2])$$

where we use the same letter ∇ for the pullback connection on h^*B .

- (b) The bracket $[\mathcal{H}, \mathcal{G}]_\nabla$: for each $\alpha_1, \alpha_2 \in \Gamma(A)$, $[\mathcal{H}, \mathcal{G}]_\nabla(\alpha_1, \alpha_2)$ is the section of the pull-back bundle $\Gamma(h^*B)$ given by:

$$(3.7) \quad [\mathcal{H}, \mathcal{G}]_\nabla(\alpha_1, \alpha_2) = -(T_\nabla(\mathcal{H}(\alpha_1), \mathcal{G}(\alpha_2)) + T_\nabla(\mathcal{G}(\alpha_1), \mathcal{H}(\alpha_2))),$$

where T_∇ is the torsion of the pullback connection on h^*B .

Now we define the **generalized Maurer-Cartan equation** for a bundle map of Lie algebroids $\mathcal{H} : A \rightarrow B$, which commutes with the anchors, to be:

$$(3.8) \quad d_\nabla \mathcal{H} + \frac{1}{2} [\mathcal{H}, \mathcal{H}]_\nabla = 0.$$

The generalized Maurer-Cartan equation is independent of the choice of connection ∇ . A fundamental fact (see [7]) is that a bundle map of Lie algebroids

¹Note that, in general, $d_\nabla^2 \neq 0$ so that d_∇ is not a differential.

$\mathcal{H} : A \rightarrow B$ which commutes with the anchors satisfies the generalized Maurer-Cartan equation if and only if it is a Lie algebroid morphism. After these preliminaries, we can turn to the proof of our result.

Proof of Proposition 3.5. First we observe that triples (M, θ, h) , where θ is a coframe on M and $h : M \rightarrow X$ is a smooth map are in 1:1 correspondence with bundle maps $\mathcal{H}_\theta = (h, \theta) : TM \rightarrow A$ which are fiberwise isomorphisms. Then one checks easily that equation (1.3) is equivalent to the condition that \mathcal{H}_θ commutes with the anchors. Moreover, if ∇ is the trivial connection on the (trivial) bundle $A = X \times \mathbb{R}^n$, then equation (1.2) is equivalent to the condition that θ be a solution of the Maurer-Cartan equation (3.8). This is a consequence of the following computation:

$$\begin{aligned}
(d_\nabla \theta + \frac{1}{2}[\theta, \theta]_\nabla)(\partial_{\theta^i}, \partial_{\theta^j}) &= \bar{\nabla}_{(\partial_{\theta^i})} \theta(\partial_{\theta^j}) - \bar{\nabla}_{(\partial_{\theta^j})} \theta(\partial_{\theta^i}) - \theta([\partial_{\theta^i}, \partial_{\theta^j}]) + \\
&\quad - T_{\bar{\nabla}}(\theta(\partial_{\theta^i}), \theta(\partial_{\theta^j})) \\
&= \nabla_{\sharp \alpha_i} \alpha_j - \nabla_{\sharp \alpha_j} \alpha_i - \theta([\partial_{\theta^i}, \partial_{\theta^j}]) + \\
&\quad - T_\nabla(\alpha_i, \alpha_j) \\
&= [\alpha_i, \alpha_j] - \sum_k \theta^k([\partial_{\theta^i}, \partial_{\theta^j}]) \alpha_k \\
&= \sum_k (C_{ij}^k - d\theta^k(\partial_{\theta^i}, \partial_{\theta^j})) \alpha_k \\
&= \sum_k \left(\sum_{l < m} C_{lm}^k \theta^l \wedge \theta^m(\partial_{\theta^i}, \partial_{\theta^j}) - d\theta^k(\partial_{\theta^i}, \partial_{\theta^j}) \right) \alpha_k
\end{aligned}$$

where $\partial_{\theta^i} = \frac{\partial}{\partial \theta^i}$ and $\bar{\nabla}$ denotes the pullback connection on h^*A . \square

Corollary 3.6. *Let $A \rightarrow X$ be the classifying Lie algebroid of a Cartan's problem. If (M, θ, h) is a realization of the problem then the rank of h is locally constant and h maps each connected component of M onto an open subset of an orbit of A .*

Proof. If $\mathcal{H}_\theta = (h, \theta) : TM \rightarrow A$ is the fiberwise isomorphism associated with the realization (M, θ, h) then the condition that it commutes with the anchors just means that:

$$dh = \sharp \circ \mathcal{H}_\theta.$$

Therefore, we conclude that for each $m_0 \in M$ (i) there is a neighborhood U of m_0 such that image $h(U_0)$ is contained in the orbit of m_0 , and (ii) $\text{rk}(d_{m_0}h)$ equals the dimension of this orbit. \square

Remark 3.7. *Note that, in general, a realization of Cartan's problem may not have constant rank. What the Corollary says is that if the problem has an associated Lie algebroid then every realization has constant rank. By Proposition 3.2, this happens if there is a solution passing through every $x_0 \in X$.*

4. MAURER-CARTAN FORMS ON LIE GROUPOIDS

For a Cartan realization problem whose manifold X reduces to a point, the structure functions C_{ij}^k are constant and the vector fields F_i vanish. In this case, the classifying algebroid is simply a Lie algebra and one can produce solutions to the realization problem by integrating this Lie algebra into a Lie group and using its Maurer-Cartan form, as was explained in Example 1.3. Our solution for a general Lie algebroid will follow the same pattern where we will now need to integrate the Lie algebroid to a Lie groupoid. Therefore, we will need to discuss first Maurer-Cartan forms on Lie groupoids and their universal property.

4.1. Definition of a Maurer-Cartan Form. In this paragraph we define Maurer-Cartan forms on Lie groupoids. These turn out to be foliated differential forms with values in the Lie algebroid.

Let \mathcal{F} be a foliation on M . Recall that an \mathcal{F} -foliated k -form on M is a section of $\wedge^k T^* \mathcal{F}$, i.e., it is a k -form on M which is only defined on k -tuples of vector fields which are all tangent to the foliation.

Definition 4.1. *An \mathcal{F} -foliated differential 1-form on M with values in a Lie algebroid $A \rightarrow X$ is a bundle map*

$$\begin{array}{ccc} T\mathcal{F} & \xrightarrow{\theta} & A \\ \downarrow & & \downarrow \\ M & \xrightarrow{h} & X \end{array}$$

which is compatible with the anchors, i.e., such that $\sharp(\theta(\xi)) = dh(\xi)$, $\forall \xi \in T\mathcal{F}$.

Let \mathcal{G} be a Lie groupoid. Right translation by an element $g \in \mathcal{G}$ is a map

$$R_g : \mathfrak{s}^{-1}(\mathfrak{t}(g)) \rightarrow \mathfrak{s}^{-1}(\mathfrak{s}(g))$$

so it does not make sense to say that a form on \mathcal{G} is right invariant. We must restrict ourselves to \mathfrak{s} -foliated differential forms. We will say that an \mathfrak{s} -foliated differential form ω on \mathcal{G} is **right invariant** if

$$\omega(\xi) = \omega((R_g)_*(\xi))$$

for every ξ tangent to an \mathfrak{s} -fiber and $g \in \mathcal{G}$. Equivalently, we will write

$$(R_g)^* \omega = \omega.$$

Remark 4.2. *If ω is a \mathfrak{s} -foliated differential form on the groupoid \mathcal{G} with values in its Lie algebroid A , the notion of right invariance still makes sense provided that we assume that $\omega : T^{\mathfrak{s}}\mathcal{G} \rightarrow A$ is a bundle map that covers the target map $\mathfrak{t} : \mathcal{G} \rightarrow X$.*

On every Lie groupoid we may define a canonical, right-invariant, differential 1-form with values in its Lie algebroid:

Definition 4.3. *The Maurer-Cartan form of a Lie groupoid \mathcal{G} is the Lie algebroid valued \mathfrak{s} -foliated 1-form*

$$\begin{array}{ccc} T^{\mathfrak{s}}\mathcal{G} & \xrightarrow{\omega_{MC}} & A \\ \downarrow & & \downarrow \\ \mathcal{G} & \xrightarrow{\mathfrak{t}} & X \end{array}$$

defined by

$$\omega_{MC}(\xi) := d_g R_{g^{-1}}(\xi) \in A_{\mathfrak{t}(g)}, \quad (\xi \in T^{\mathfrak{s}}_g \mathcal{G}).$$

For any foliation \mathcal{F} of M , the bundle $T\mathcal{F} \rightarrow M$ has an obvious Lie algebroid structure with anchor the inclusion. Hence, it makes sense to talk about the generalized Maurer-Cartan equation in this context (see Section 3.2), and we have the following:

Proposition 4.4. *The Maurer-Cartan form ω_{MC} of \mathcal{G} satisfies the generalized Maurer-Cartan equation. Hence, we can view the Maurer-Cartan form as a Lie algebroid morphism $\omega_{MC} : T^{\mathfrak{s}}\mathcal{G} \rightarrow A$.*

Proof. This proposition is a consequence of the following calculation:

$$\begin{aligned}
 d_{\nabla}\omega_{\text{MC}}(\xi_1, \xi_2) &= \nabla_{\mathbf{t}_*\xi_1}\omega_{\text{MC}}(\xi_2) - \nabla_{\mathbf{t}_*\xi_2}\omega_{\text{MC}}(\xi_1) - \omega_{\text{MC}}([\xi_1, \xi_2]) \\
 &= \nabla_{\#}\alpha\beta - \nabla_{\#}\beta\alpha - [\alpha, \beta] \\
 &= T_{\nabla}(\alpha, \beta) \\
 &= T_{\nabla}(\omega_{\text{MC}}(\xi_1), \omega_{\text{MC}}(\xi_2)) = -\frac{1}{2}[\omega_{\text{MC}}, \omega_{\text{MC}}](\xi_1, \xi_2).
 \end{aligned}$$

where ξ_1 and ξ_2 are right invariant vector fields on \mathcal{G} and $\alpha, \beta \in \Gamma(A)$ are their generators. \square

4.2. The Local Universal Property. We will now show that any 1-form on a differentiable manifold, with values in an integrable Lie algebroid A , satisfying the generalized Maurer-Cartan equation is locally the pullback of the Maurer-Cartan form on a Lie groupoid integrating A . We will need the following lemma.

Lemma 4.5. *Let \mathcal{F} be a foliation on a manifold M and let θ be an \mathcal{F} -foliated 1-form (over h) on M with values in a Lie algebroid $A \rightarrow X$ equipped with an arbitrary connection ∇ . Assume that the distribution $\mathcal{D} = \{\ker \theta_x : x \in M\} \subset T\mathcal{F} \subset TM$ has constant rank. Then \mathcal{D} is integrable if and only if $d_{\nabla}\theta(\xi_1, \xi_2) = 0$ whenever $\xi_1, \xi_2 \in \mathcal{D}$.*

Proof. Choose a local basis $\xi_1, \dots, \xi_r \in \mathfrak{X}(\mathcal{F})$ of \mathcal{D} in an open set of M . Then by Frobenius Theorem, \mathcal{D} is integrable if and only if $[\xi_i, \xi_j] \in \text{span}\{\xi_1, \dots, \xi_r\}$ for all $1 \leq i, j \leq r$, which happens if and only if $\theta([\xi_i, \xi_j]) = 0$ for all $1 \leq i, j \leq r$. Since $\theta(\xi_i) = 0$ for all $1 \leq i \leq r$ we have

$$(4.1) \quad d_{\nabla}\theta(\xi_i, \xi_j) = \bar{\nabla}_{\xi_i}\theta(\xi_j) - \bar{\nabla}_{\xi_j}\theta(\xi_i) - \theta([\xi_i, \xi_j]) = -\theta([\xi_i, \xi_j])$$

from which the result follows. \square

Theorem 4.6 (Local Universal Property). *Let θ be a 1-form (over h) on a manifold M , with values in an integrable Lie algebroid A , that satisfies the generalized Maurer-Cartan equation. Let \mathcal{G} be a Lie groupoid integrating A and denote by ω_{MC} its right invariant Maurer-Cartan form. Then, for each $p_0 \in M$ and $g_0 \in \mathcal{G}$ such that $h(p_0) = \mathbf{t}(g_0)$, there exists a unique locally defined diffeomorphism $\phi : M \rightarrow \mathbf{s}^{-1}(h(p_0))$ satisfying*

$$\phi(p_0) = g_0, \quad \phi^*\omega_{\text{MC}} = \theta.$$

Remark 4.7. *We can summarize the theorem by saying that, at least locally, there is a unique map $\phi : M \rightarrow \mathcal{G}$ which makes the following diagram of Lie algebroid morphisms commute:*

$$\begin{array}{ccc}
 TM & \overset{\phi_*}{\dashrightarrow} & T^s\mathcal{G} \\
 \downarrow \theta & & \swarrow \omega_{\text{MC}} \\
 & & A \\
 & & \downarrow \phi \\
 M & \overset{\phi}{\dashrightarrow} & \mathcal{G} \\
 \downarrow h & & \swarrow \mathbf{t} \\
 & & X
 \end{array}$$

Proof. The proof uses the usual technique of the graph, so we will construct the graph of ϕ by integrating a convenient distribution. Uniqueness will then follow from the uniqueness of integral submanifolds of a distribution. Let us set:

$$M_{h \times \mathbf{t}} \mathcal{G} = \{(p, g) \in M \times \mathcal{G} : h(p) = \mathbf{t}(g)\}.$$

Since \mathbf{t} is a surjective submersion, this fibered product over X is a manifold and it comes equipped with the foliation \mathcal{F} given by the fibers of $\mathbf{s} \circ \pi_{\mathcal{G}}$. On $M_{h \times \mathbf{t}} \mathcal{G}$ consider the A -valued \mathcal{F} -foliated 1-form

$$\Omega = \pi_M^* \theta - \pi_{\mathcal{G}}^* \omega_{\text{MC}}.$$

Let $\mathcal{D} = \ker \Omega$ denote the associated distribution on $M_{h \times \mathbf{t}} \mathcal{G}$.

In order to apply the previous lemma, we must first show that \mathcal{D} has constant rank. We will do this by showing that

$$(d\pi_M)_{(p,g)}|_{\mathcal{D}_{(p,g)}} : \mathcal{D}_{(p,g)} \rightarrow T_p M$$

is an isomorphism for each $(p, g) \in M_{h \times \mathbf{t}} \mathcal{G}$. Note that this also implies that if \mathcal{D} is integrable then its leaf through (p_0, g_0) is locally the graph of a uniquely defined diffeomorphism $\phi : M \rightarrow \mathbf{s}^{-1}(\mathbf{t}(g_0))$ satisfying $\phi(p_0) = g_0$.

Suppose that $(d\pi_M)_{(p,g)}(v, w) = 0$ for some $(v, w) \in \mathcal{D}_{(p,g)}$. Since \mathcal{D} is contained in $T\mathcal{F}$, it follows that w is \mathbf{s} -vertical and $\omega_{\text{MC}}(w)$ is simply the right translation of w to $\mathbf{1}_{\mathbf{t}(g)}$ and thus,

$$\begin{aligned} (d\pi_M)_{(p,g)}(v, w) = 0 &\implies v = 0 \\ &\implies \omega_{\text{MC}}(w) = 0 \quad (= \theta(v)) \\ &\implies w = 0 \\ &\implies (v, w) = 0 \end{aligned}$$

so that $(d\pi_M)_{(p,g)}$ is injective. Now, if $v \in T_p M$ then $(v, (R_g)_* \theta(v))$ is an element of $\mathcal{D}_{(p,g)}$ so $(d\pi_M)_{(p,g)}$ is also surjective.

Having accomplished this, we may use the preceding lemma to complete the proof. We compute (omitting the pullbacks for simplicity):

$$\begin{aligned} d_{\nabla} \Omega &= d_{\nabla} \theta - d_{\nabla} \omega_{\text{MC}} \\ &= -\frac{1}{2}[\theta, \theta] + \frac{1}{2}[\omega_{\text{MC}}, \omega_{\text{MC}}]. \end{aligned}$$

Replacing $\theta = \Omega + \omega_{\text{MC}}$ we obtain

$$\begin{aligned} d_{\nabla} \Omega &= -\frac{1}{2}[\Omega + \omega_{\text{MC}}, \Omega + \omega_{\text{MC}}] + \frac{1}{2}[\omega_{\text{MC}}, \omega_{\text{MC}}] \\ &= -\frac{1}{2}[\Omega, \Omega] - \frac{1}{2}[\Omega, \omega_{\text{MC}}] - \frac{1}{2}[\omega_{\text{MC}}, \Omega]. \end{aligned}$$

So $d_{\nabla} \Omega(\xi_1, \xi_2) = 0$ whenever $\Omega(\xi_1) = 0 = \Omega(\xi_2)$. Hence \mathcal{D} is integrable and the proof is completed. \square

Remark 4.8. *With a slight modification, the theorem above is still valid even when A is not integrable. In fact, since an A -valued Maurer-Cartan form on M*

$$(\theta, h) \in \Omega^1(M, A)$$

is the same as a Lie algebroid morphism

$$\begin{array}{ccc} TM & \xrightarrow{\theta} & A \\ \downarrow & & \downarrow \\ M & \xrightarrow{h} & X \end{array}$$

it follows that $h(M)$ lies in a single orbit of A in X . By restricting h to a small enough neighborhood, we may assume that its image is a contractible open set $U \subset L$ in an orbit L of A in X . Then, the restriction of A to U is integrable [5] and we may proceed as in the proof of the theorem.

As a consequence of the theorem we obtain the following useful corollary:

Corollary 4.9. *Let \mathcal{G} be a Lie groupoid with Maurer-Cartan form ω_{MC} and let $x, y \in X$ be points in the same orbit. If $\phi : \mathfrak{s}^{-1}(x) \rightarrow \mathfrak{s}^{-1}(y)$ is a symmetry of ω_{MC} (i.e., $\phi^*\omega_{MC} = \omega_{MC}$) then $\phi = R_g$ for some $g \in \mathcal{G}$.*

Proof. Note that the equation $\phi^*\omega_{MC} = \omega_{MC}$ only makes sense provided $\mathfrak{t} \circ \phi = \phi$ (see Remark 4.2). Therefore, $g := \phi(\mathbf{1}_x)$ is an arrow joining y to x and

$$R_g : \mathfrak{s}^{-1}(x) \rightarrow \mathfrak{s}^{-1}(y), h \mapsto hg$$

is a symmetry of ω_{MC} . Since $R_g\mathbf{1}_x = g = \phi(\mathbf{1}_x)$, from the uniqueness part of Theorem 4.6 we must have $\phi = R_g$. \square

4.3. The Global Universal Property. There is a more conceptual proof of Theorem 4.6 that will lead us to a global version of the universal property of Maurer-Cartan forms.

First, observe that Theorem 4.6 is a local result so we may assume that M is simply connected. The source simply connected Lie groupoid integrating TM is then the pair groupoid $M \times M \rightrightarrows M$. By Lie's Second Theorem (see [7]), there exists a unique morphism of Lie groupoids

$$\begin{array}{ccc} M \times M & \xrightarrow{H} & \mathcal{G} \\ \Downarrow & & \Downarrow \\ M & \xrightarrow{h} & X \end{array}$$

integrating θ . Now, if we are given $p_0 \in M$ and $g_0 \in \mathcal{G}$ with $h(p_0) = \mathfrak{s}(g_0)$, then we may write

$$H(p, p') = \phi(p)\phi(p')^{-1}$$

where $\phi : M \rightarrow \mathfrak{s}^{-1}(h(p_0)) \subset \mathcal{G}$ is defined by $\phi(p) := H(p, p_0)g_0$. Note that ϕ satisfies

$$\phi(p_0) = g_0, \quad \phi^*\omega_{MC} = \theta,$$

and so it has the desired properties.

In general, when M is not simply connected, the Lie algebroid morphism θ integrates to a Lie groupoid morphism

$$\begin{array}{ccc} \Pi_1(M) & \xrightarrow{F} & \mathcal{G} \\ \Downarrow & & \Downarrow \\ M & \xrightarrow{h} & X \end{array}$$

where $\Pi_1(M)$ denotes the fundamental groupoid of M . The problem of determining when θ is globally the pullback of the Maurer-Cartan form on a Lie groupoid \mathcal{G} integrating A is then reduced to determining when the morphism F above factors through the groupoid covering projection

$$\begin{aligned} p : \Pi_1(M) &\rightarrow M \times M, \\ p([\gamma]) &:= (\gamma(1), \gamma(0)). \end{aligned}$$

Theorem 4.10 (Global Universal Property). *Let A be an integrable Lie algebroid with source simply connected Lie groupoid $\mathcal{G}(A)$ and $(\theta, h) \in \Omega^1(M, A)$ be an A -valued differential 1-form on M . Then, there exists a globally defined local diffeomorphism*

$$\phi : M \rightarrow \mathfrak{s}^{-1}(h(p_0)), \quad \phi(p_0) = g_0, \quad \phi^* \omega_{MC} = \theta,$$

if and only if

Local obstruction:: *The 1-form θ satisfies the generalized Maurer-Cartan equation and*

Global obstruction:: *The morphism F integrating θ is trivial when restricted to the fundamental group $\pi_1(M, p_0)$ (i.e., the isotropy group at p_0).*

Proof. We begin by proving that both conditions are necessary. It is clear that if $\theta = \phi^* \omega_{MC}$ then θ satisfies the generalized Maurer-Cartan equation. So all we have to prove is that F is trivial on the isotropy group $\pi_1(M, p_0)$. Suppose that ϕ exists. Then the map

$$H : M \times M \rightarrow \mathcal{G}$$

given by

$$H(p, p') = \phi(p)\phi(p')^{-1}.$$

defines a Lie groupoid morphism over the map $\mathfrak{t} \circ \phi$.

It then follows from $\phi^* \omega_{MC} = \theta$ that $\mathfrak{t} \circ \phi = h$ and that H integrates θ . To see this, notice that if $f : M \rightarrow \mathfrak{s}^{-1}(x)$ and $g : M \rightarrow \mathfrak{t}^{-1}(x)$ are smooth maps, then the differential of $\varphi(p, p') = f(p) \cdot g(p')$ is given by

$$(\mathrm{d}\varphi)_{(p,p')}(v, w) = (\mathrm{d}L_{f(p)})_{g(p')}(\mathrm{d}g)_{p'}(w) + (\mathrm{d}R_{g(p')})_{f(p)}(\mathrm{d}f)_p(v)$$

for $v, w \in T_{(p,p')}(M \times M)$. Thus, in our case we obtain, for any $v \in T_p M$,

$$\begin{aligned} (\mathrm{d}H)_{(p,p)}(0, v) &= (\mathrm{d}R_{\phi(p)^{-1}})_{\phi(p)}(\mathrm{d}\phi)_p(v) \\ &= \omega_{MC}(\phi_* v) \\ &= \phi^* \omega_{MC}(v) \\ &= \theta(v). \end{aligned}$$

Finally, since H integrates θ we obtain the commutative diagram

$$\begin{array}{ccc} \Pi_1(M) & \xrightarrow{F} & \mathcal{G} \\ p \downarrow & \nearrow H & \\ M \times M & & \end{array}$$

where p denotes the covering projection $p([\gamma]) = (\gamma(1), \gamma(0))$. We conclude that F only depends on the endpoints of γ and not on its homotopy class, proving the claim.

Conversely, suppose both conditions are satisfied. Then by Theorem 4.6, it follows that θ is locally the pullback of ω_{MC} by a map ϕ_{loc} . However, since F only depends on the end points of paths γ and not on their homotopy class, F factors through

$$\begin{array}{ccc} \Pi_1(M) & \xrightarrow{F} & \mathcal{G} \\ p \downarrow & \dashrightarrow H & \\ M \times M & & \end{array}$$

Thus, by setting

$$\phi(p) := H(p, p_0)g_0$$

we obtain a global map which, by the uniqueness result in Theorem 4.6, restricts to the locally defined maps ϕ_{loc} . It follows that $\phi^*\omega_{\text{MC}} = \theta$ and $\phi(p_0) = g_0$ proving the theorem. \square

An immediate consequence of the global universal property is the following corollary:

Corollary 4.11. *Let U be a contractible manifold, let $A \rightarrow X$ be a weakly integrable Lie algebroid and let $h : U \rightarrow X$ be a smooth map. Any two A -valued 1-forms $\theta_1, \theta_2 \in \Omega^1(U; A)$ over h satisfying the generalized Maurer-Cartan equation are equivalent: for any $p \in U$ there exists a diffeomorphism $\phi : U \rightarrow U$, fixing p and commuting with h , such that $\phi^*\theta_2 = \theta_1$.*

Remark 4.12. *Using the A -path approach to integrability, introduced in [5], one can express the global obstruction condition of the preceding theorem at the infinitesimal level, i.e., without any mention to the Lie groupoid \mathcal{G} integrating A . In fact, given any curve $\gamma : I \rightarrow M$, the path $\theta \circ \dot{\gamma} : I \rightarrow A$ satisfies*

$$\sharp(\theta \circ \dot{\gamma}(t)) = \frac{d}{dt}(h \circ \gamma)(t) \text{ for all } t \in I$$

and thus, is an A -path. We can then rewrite the condition as:

Global obstruction: For every loop γ in M , homotopic to the constant curve at p , the A -path $\theta \circ \dot{\gamma}$ is A -homotopic to the constant zero A -path at $h(p)$.

Note also that this last condition can be expressed in terms of a differential equation (see [5]).

5. LOCAL CLASSIFICATION AND SYMMETRIES

We now return to Cartan's realization problem. We will use the Lie groupoid integrating the classifying Lie algebroid to solve the Local Classification Problem and to study the symmetries of the problem.

5.1. Local Classification Problem. The results of the previous two sections lead to a solution to the Local Classification Problem. Before we give the main result, we must say a few words about equivalence of realizations. By this we mean:

Definition 5.1. *Let (n, X, C_{ij}^k, F_i) be the initial data of a Cartan's realization. Given two realizations (M_1, θ_1, h_1) and (M_2, θ_2, h_2) a (locally defined) diffeomorphism $\phi : M_1 \rightarrow M_2$ is called a (local) **equivalence of realizations** if*

$$\phi^*\theta_2^i = \theta_1^i, \quad h_2 \circ \phi = h_1.$$

This notion of equivalence should not be confused with the notion of equivalence of coframe.

Remark 5.2. *Note that if θ is a coframe on M and $A_\theta \rightarrow X_\theta$ is the corresponding classifying Lie algebroid, then we have the realization $(M, \theta, \kappa_\theta)$. In this case, a self equivalence of the realization $\phi : (M, \theta, \kappa_\theta) \rightarrow (M, \theta, \kappa_\theta)$ is the same thing as an equivalence of the coframe $\phi : (M, \theta) \rightarrow (M, \theta)$, because any self-equivalence ϕ of the coframe must satisfy $\kappa_\theta \circ \phi = \kappa_\theta$. Also, given two realizations (M_1, θ_1, h_1) and (M_2, θ_2, h_2) , any equivalence of coframes $\phi : (M_1, \theta_1) \rightarrow (M_2, \theta_2)$ obviously determines an equivalence of realizations $\phi : (M_1, \theta_1, h_2 \circ \phi) \rightarrow (M_2, \theta_2, h_2)$ but not necessarily of the original realizations, since the identity $h_2 \circ \phi = h_1$ may not hold.*

Now we can state:

Theorem 5.3 (Local Classification). *Let (n, X, C_{ij}^k, F_i) be the initial data of a Cartan's realization problem and denote by $A \rightarrow X$ its classifying Lie algebroid. Then:*

- (i) *Any realization is locally equivalent to a neighborhood of the identity of a fiber $\mathbf{s}^{-1}(x_0)$ of a groupoid \mathcal{G} integrating A , equipped with the Maurer-Cartan form.*
- (ii) *Two realizations are locally equivalent if and only if they correspond to the same point $x_0 \in X$.*

Proof. Recall that we have a basis $\{\alpha_1, \dots, \alpha_n\}$ of global sections of the classifying Lie algebroid $A \rightarrow X$ such that:

$$[\alpha_i, \alpha_j] = -C_{ij}^k \alpha_k, \quad \sharp \alpha_i = F_i.$$

Let us suppose for the moment that A is an integrable Lie algebroid and that \mathcal{G} is a Lie groupoid integrating A . We denote by ω_{MC} the Maurer-Cartan form of \mathcal{G} and by $(\omega_{\text{MC}}^1, \dots, \omega_{\text{MC}}^n)$ its components with respect to the basis $\{\alpha_1, \dots, \alpha_n\}$. Then, for each $x_0 \in X$, $(\mathbf{s}^{-1}(x_0), \omega_{\text{MC}}^i, \mathbf{t})$ is a realization of Cartan's problem with initial data (n, X, C_{ij}^k, F_i) .

A similar argument still holds when A is not integrable. In this case, for each $x_0 \in X$ we can find a neighborhood $U \subset L$ of x_0 in the leaf L containing it, such that the restriction of A to U is integrable to a Lie groupoid $\mathcal{G} \rightrightarrows U$. The Maurer-Cartan form of \mathcal{G} takes values in $A|_U \hookrightarrow A$ so we can see it as an A -valued Maurer-Cartan form. It is again clear that $(\mathbf{s}^{-1}(x_0), \omega_{\text{MC}}^i, \mathbf{t})$ is a realization of (n, X, C_{ij}^k, F_i) .

If (M, θ^i, h) is any realization of (n, X, C_{ij}^k, F_i) , Proposition 3.5 shows that the A -valued 1-form $\theta \in \Omega^1(M, A)$ defined by

$$\theta = \sum_i \theta^i(\alpha_i \circ h)$$

satisfies the generalized Maurer-Cartan equation.

$$d_{\nabla} \theta + \frac{1}{2}[\theta, \theta]_{\nabla} = 0.$$

Therefore, if $p_0 \in M$ is such that $h(p_0) = x_0$ then, by the universal property of Maurer-Cartan forms, we can find a neighborhood V of p_0 in M and a unique diffeomorphism $\phi : V \rightarrow \phi(V) \subset \mathbf{s}^{-1}(x_0)$ such that $\phi(p_0) = \mathbf{1}_{x_0}$ and $\phi^* \omega_{\text{MC}} = \theta$. This means that any realization of Cartan's problem is locally equivalent to a neighborhood of the identity of an \mathbf{s} -fiber of \mathcal{G} equipped with the Maurer-Cartan form.

Finally, if (M_1, θ_1, h_1) and (M_2, θ_2, h_2) are two realizations of (n, X, C_{ij}^k, F_i) and $h_1(p_1) = h_2(p_2) = x_0$, then there exist open sets $V_i \subset M_i$ and (unique) diffeomorphisms $\phi_i : V_i \rightarrow \phi_i(V_i) \subset \mathbf{s}^{-1}(x_0)$ such that $\phi_i(p_i) = \mathbf{1}_{x_0}$ and $\phi_i^* \omega_{\text{MC}} = \theta_i$, ($i = 1, 2$). Hence, the map $\phi_2^{-1} \circ \phi_1$ is a local equivalence from $\phi_1^{-1}(\phi_1(V_1) \cap \phi_2(V_2)) \subset M_1$ onto $\phi_2^{-1}(\phi_1(V_1) \cap \phi_2(V_2)) \subset M_2$. The converse is obvious, since by definition an equivalence between two realizations (M_1, θ_1, h_1) and (M_2, θ_2, h_2) of (n, X, C_{ij}^k, F_i) must satisfy $h_2 \circ \phi = h_1$. \square

5.2. Equivalence versus formal equivalence. Another consequence of the universal property of the Maurer-Cartan form is that “equivalence” and “formal equivalence” for a coframe actually coincide.

Proposition 5.4. *Let θ be a fully regular coframe on a manifold M . Any two points $p, q \in M$ are equivalent if and only if they are formally equivalent.*

Proof. Clearly, if two points are equivalent then they are formally equivalent. For the converse, let $\phi : U \rightarrow V$ be a formal equivalence defined on contractible open

sets such that $\phi(p) = q$. Denote by $A_\theta \rightarrow X_\theta$ the classifying Lie algebroid of the coframe and by $\kappa_\theta : M \rightarrow X_\theta$ the classifying map. Then, by its very definition, we find that $\kappa_\theta(U) = \kappa_\theta(V) = W$ is an open set of X_θ and that the following diagram commutes:

$$\begin{array}{ccc} U & \xrightarrow{\phi} & V \\ & \searrow \kappa_\theta & \swarrow \kappa_\theta \\ & & W \end{array}$$

Our aim is to prove that there exists a diffeomorphism $\psi : U \rightarrow V$ such that $\psi(p) = q$ and $\psi^*\theta|_V = \theta|_U$.

For this, observe that since ϕ is a formal equivalence we have $C_{ij}^k = C_{ij}^k \circ \phi$, so the coframe $\phi^*(\theta|_V)$ satisfies the same structure equations as $\theta|_U$:

$$d(\phi^*\theta^k) = \sum_{i < j} C_{ij}^k(\phi^*\theta^i) \wedge (\phi^*\theta^j).$$

It follows that we have two Lie algebroid morphisms:

$$\begin{array}{ccc} TU & \xrightarrow{\phi^*\theta} & A_\theta|_W \\ \downarrow & & \downarrow \\ U & \xrightarrow{\kappa_\theta} & W \end{array} \quad \begin{array}{ccc} TU & \xrightarrow{\theta} & A_\theta|_W \\ \downarrow & & \downarrow \\ U & \xrightarrow{\kappa_\theta} & W \end{array}$$

We conclude that $\phi^*(\theta|_V)$ and $\theta|_U$ are both 1-forms in $\Omega^1(U, A_\theta)$ over κ_θ satisfying the Maurer-Cartan equation. By Corollary 4.11, we conclude that there exists a diffeomorphism $\varphi : U \rightarrow U$, fixing $p \in U$ and commuting with κ_θ , such that $\varphi^*(\phi^*(\theta|_V)) = \theta|_U$. The map $\psi := \phi \circ \varphi : U \rightarrow V$ is the desired equivalence mapping p to q . \square

5.3. Symmetries of Realizations. We can also use our solution to the Local Equivalence Problem to recover a few classical results about symmetries of coframes. Many of these results can be traced back to Cartan [3]. The formulation presented here is based on [1]. Note that our only purpose is to show how the classifying Lie algebroid can be used to give very simple proofs of these facts.

Definition 5.5. *Let θ be a coframe on M . A **symmetry** of (M, θ) is a self-equivalence, i.e., a diffeomorphism $\phi : M \rightarrow M$ such that $\phi^*\theta = \theta$. An **infinitesimal symmetry** is a vector field $\xi \in \mathfrak{X}(M)$ such that $\mathcal{L}_\xi\theta = 0$.*

Clearly, the set of infinitesimal symmetries of a coframe (M, θ) form a Lie subalgebra $\mathfrak{X}(M, \theta) \subset \mathfrak{X}(M)$. On the other hand, we recall that the group of diffeomorphism $\text{Diff}(M)$ is a Fréchet Lie group for the compact-open C^∞ -topology. The set of all symmetries of a coframe (M, θ) form a topological subgroup $\text{Diff}(M, \theta) \subset \text{Diff}(M)$ for the induced topology.

We also have the related notions of symmetry and infinitesimal symmetry of a realization:

Definition 5.6. *If (M, θ, h) is a realization of Cartan's problem we call a diffeomorphism $\phi : M \rightarrow M$ a **symmetry of the realization** if $\phi^*\theta = \theta$ and $h \circ \phi = h$. Similarly, one calls a vector field $\xi \in \mathfrak{X}(M)$ an **infinitesimal symmetry of the realization** if $\mathcal{L}_\xi\theta = 0$ and $dh \cdot \xi = 0$.*

Given a realization (M, θ, h) we will denote by $\text{Diff}(M, \theta, h)$ the group of symmetries of the realization and by $\mathfrak{X}(M, \theta, h)$ the Lie algebra of infinitesimal symmetries of the realization. Obviously, $\text{Diff}(M, \theta, h)$ is a subgroup of the group $\text{Diff}(M, \theta)$ of

symmetries of the underlying coframe, while $\mathfrak{X}(M, \theta, h)$ is a Lie subalgebra of the Lie algebra $\mathfrak{X}(M, \theta)$ of infinitesimal symmetries of the underlying coframe.

Our next result describes the relationship between the symmetries and the classifying Lie algebroid:

Proposition 5.7 (Theorem A.2 of [1]). *Let (M, θ, h) be a realization of a Cartan's problem with classifying Lie algebroid $A \rightarrow X$. Then the set $\mathfrak{X}(M, \theta, h)_p$ of germs of infinitesimal symmetries of (M, θ, h) at a point p is a Lie algebra isomorphic to $\mathfrak{g}_{h(p)}$, the isotropy Lie algebra of A at $h(p)$. In particular, if M is connected then $h(M)$ is an open subset of a leaf L of A and*

$$\dim \mathfrak{X}(M, \theta, h)_p = \dim M - \dim L$$

Proof. Since the result is local, we may assume without loss of generality that the classifying Lie algebroid $A \rightarrow X$ is integrable. If this is not the case, we can restrict A to a contractible open set U in the leaf containing $h(p)$ so that $A|_U$ is integrable. Let $\mathcal{G} \rightrightarrows X$ be a Lie groupoid with Lie algebroid A . Then, by the local universal property of the Maurer-Cartan form on \mathcal{G} (Theorem 4.6), there is a neighborhood U of p in M and a diffeomorphism $\phi : U \rightarrow \phi(U) \subset \mathfrak{s}^{-1}(h(p))$ such that $\phi(p) = \mathbf{1}_{h(p)}$ and $\phi^* \omega_{\text{MC}} = \theta$.

It follows that $\mathfrak{X}(M, \theta, h)_p$ coincides with the set of germs at $\mathbf{1}_{h(p)}$ of vector fields tangent to $\mathfrak{s}^{-1}(h(p))$ which are infinitesimal symmetries of the Maurer-Cartan form. Thus, from (the infinitesimal version of) Corollary 4.9 we may conclude that these may be identified with the isotropy Lie algebra of A at p . \square

The previous result applies, in particular, to the classifying algebroid A_θ of the coframe. Hence, if M is connected, we see that for any $p, q \in M$ the Lie algebras $\mathfrak{X}(M, \theta)_p$ and $\mathfrak{X}(M, \theta)_q$ are isomorphic. In the case of coframes, the following result is classical (see Theorem 3.2 of [11]):

Theorem 5.8. *Let (M, θ, h) be a realization of a Cartan's problem. Then its group of symmetries $\text{Diff}(M, \theta, h)$ is a (finite dimensional) Lie group of transformations of M with Lie algebra $\mathfrak{X}_c(M, \theta, h) \subset \mathfrak{X}(M, \theta, h)$ the subspace generated by the complete infinitesimal symmetries of the realization.*

Proof. We can assume that M is connected. A classical theorem of Palais (Theorem IV.III of [15]) states that any group $G \subset \text{Diff}(M)$ such that the Lie subalgebra $\mathfrak{g} \subset \mathfrak{X}(M)$ generated by the complete vector fields X whose flows $\phi_X^t \in G$ is finite dimensional, is actually a Lie group of transformations of M with Lie algebra \mathfrak{g} . Hence, by Proposition 5.7, it is enough to prove that for any $p \in M$ the restriction map $\mathfrak{X}(M, \theta, h) \rightarrow \mathfrak{X}(M, \theta, h)_p$ is injective. This follows from the following lemma:

Lemma 5.9. *Let $X \in \mathfrak{X}(M, \theta, h)$ be an infinitesimal symmetry and assume that X vanishes at some point $p \in M$. Then $X \equiv 0$.*

To prove this lemma, it is enough to prove that the zero set of $X \in \mathfrak{X}(M, \theta, h)$ is both open and closed. It is obviously closed, and to prove that it is open let $q \in M$ be such that $X_q = 0$. If $\{X_i\}$ are the linearly independent vector fields dual to the coframe $\{\theta^i\}$, then:

$$\mathcal{L}_X \theta^i = 0 \quad \Leftrightarrow \quad \mathcal{L}_X X_i = 0.$$

Hence, we have that:

$$X_q = 0 \text{ and } \mathcal{L}_{X_i} X = 0.$$

Since the X_i are everywhere linearly independent, we conclude that X vanishes in a neighborhood of q . \square

We also have the following semi-global symmetry property of a realization:

Corollary 5.10 (Theorem A.3 of [1]). *Let (n, X, C_{ij}^k, F_i) be the initial data of a realization problem with associated Lie algebroid $A \rightarrow X$. Let $L \subset X$ be a leaf of A whose isotropy Lie algebra is \mathfrak{g} , and let G be any Lie group with Lie algebra \mathfrak{g} . Then over any contractible set $U \subset L$, there exists a principal G -bundle $h : M \rightarrow U$ and a G -invariant coframe θ of M such that (M, θ, h) is a realization of the Cartan's problem. Moreover, this realization is locally unique up to isomorphism.*

Proof. This is an immediate consequence of our classification result, Theorem 5.3. In fact, since U is contractible and the restriction of A to U is transitive, it follows that $A|_U$ is integrable. Moreover, for any Lie group G with Lie algebra \mathfrak{g} , there exists a Lie groupoid \mathcal{G} integrating $A|_U$ whose isotropy Lie group is isomorphic to G . The restriction of the Maurer-Cartan form of \mathcal{G} to any \mathfrak{s} -fiber furnishes the desired (locally unique) G -invariant coframe. \square

We can also use Theorem 5.3 to explain the relationship between the classifying algebroid $A \rightarrow X$ of a Cartan realization problem and the classifying algebroid of a coframe θ associated with a realization (M, θ, h) :

Corollary 5.11. *Let (n, X, C_{ij}^k, F_i) be a Cartan realization problem with associated classifying algebroid A and let (M, θ, h) be a connected realization. Then $h(M)$ is an open subset of an orbit of A and there is a Lie algebroid morphism from the Lie subalgebroid $A|_{h(M)} \subset A$ to $A_\theta \rightarrow X_\theta$ (the classifying algebroid of the coframe θ) which is a fiberwise isomorphism that covers a surjective submersion.*

Proof. Since M is connected, $h(M)$ is an open subset of an orbit of A . Also, we have a two leg diagram:

$$\begin{array}{ccc} & M & \\ h \swarrow & & \searrow \kappa_\theta \\ h(M) & \overset{\psi}{\dashrightarrow} & X_\theta \end{array}$$

Observe that if $h(p) = h(q)$ then Theorem 5.3 gives a locally defined diffeomorphism $\phi : M \rightarrow M$, such that $\phi(p) = q$ and $\phi^*\theta = \theta$. In other words, p and q are locally formally equivalent so that we must have $\kappa_\theta(p) = \kappa_\theta(q)$. This shows that we have a well defined map $\psi : h(M) \rightarrow X_\theta$ that makes the diagram above commutative. Since κ_θ is a surjective submersion, we conclude that $\psi : h(M) \rightarrow X_\theta$ is also a surjective submersion.

Finally, observe that the coframe yields Lie algebroid maps

$$\begin{array}{ccc} & TM & \\ \swarrow & & \searrow \\ A|_{h(M)} & \dashrightarrow & A_\theta \end{array}$$

which cover the two legs in the diagram above, and which are fiberwise isomorphism. Hence, we conclude that there is a Lie algebroid morphism $A|_{h(M)} \rightarrow A_\theta$, over the map $\psi : h(M) \rightarrow X_\theta$, which is a fiberwise isomorphism. \square

Remark 5.12. *The corollary above can be interpreted as saying that the realization problem associated to a coframe is the one that has the least amount of invariant functions involved (and thus the largest symmetry group). On the other hand, a general realization problem deals with a coframe along with extra invariants that must be preserved.*

The following example illustrates the point made in this remark.

Example 5.13. Suppose that $\Psi : G \times X \rightarrow X$ is an action of an n -dimensional Lie group G on a manifold X , and let $\psi : \mathfrak{g} \rightarrow \mathfrak{X}(X)$ be the associated infinitesimal action. Let us denote by $A = \mathfrak{g} \times X$ the corresponding transformation Lie algebroid. Since A is a trivial vector bundle, it determines a realization problem. In fact, if e_1, \dots, e_n is a basis of \mathfrak{g} such that

$$[e_i, e_j] = C_{ij}^k e_k \text{ and } F_i = \psi(e_i),$$

then (n, X, C_{ij}^k, F_i) is the initial data to a realization whose classifying Lie algebroid is A .

It then follows from Theorem 5.3 that every solution to this realization problem is locally equivalent to a neighborhood of the identity in $(G, \omega_{MC}^i, \Psi_x)$, where ω_{MC}^i are the components of the Maurer-Cartan form on G with respect to the basis e_1, \dots, e_n of \mathfrak{g} and

$$\Psi_x : G \rightarrow X, \quad \Psi_x(g) = \Psi(g, x).$$

Thus, the infinitesimal symmetry algebra of a realization that corresponds to a point $x \in X$ is the isotropy Lie algebra of the infinitesimal action, i.e., $\mathfrak{X}(G, \omega_{MC}^i, \Psi_x)|_{\text{Id}} = \ker \psi_x$. Moreover, a map $\phi : G \rightarrow G$ which preserves the Maurer-Cartan form is an equivalence, if and only if it leaves the action of G on the orbit through x invariant. In other words, it must be the right translation by an element of the isotropy subgroup G_x .

On the other hand, the classifying Lie algebroid of the Maurer-Cartan coframe on G is the Lie algebra \mathfrak{g} itself. The symmetries of the Maurer-Cartan coframe on G are the right translations by any element of G .

6. GLOBAL EQUIVALENCE

We now turn to global questions related to equivalence of coframes. Namely, we will give our solution to the Globalization and to the Global Equivalence Problems

6.1. The Globalization Problem. Recall that the *globalization problem* (see Problem 1.8) asks if two germs of coframes θ_0 and θ_1 which solve the same realization problem are germs of the same global realization.

We recall that we say that a Lie algebroid A is *weakly integrable* if the restriction of A to any orbit is integrable. Then we have:

Theorem 6.1. Consider a Cartan problem whose associated classifying Lie algebroid $A \rightarrow X$ is weakly integrable. Then (M_0, θ_0, h_0) and (M_1, θ_1, h_1) are germs of the same global connected realization (M, θ, h) if and only if they correspond to points on X in the same orbit of A .

Proof. Given the two germs (M_0, θ_0, h_0) and (M_1, θ_1, h_1) , suppose that they correspond to points x_0 and x_1 on the same orbit L of A : $h_0(m_0) = x_0$ and $h_1(m_1) = x_1$. Since A is weakly integrable, there exists a Lie groupoid \mathcal{G}_L integrating $A|_L$ and the \mathfrak{s} -fiber at x_0 contains a point g with $\mathfrak{t}(g) = x_1$. Thus, the germ of (M_0, θ_0, h_0) at m_0 can be identified with the germ of $(\mathfrak{s}^{-1}(x_0), \omega_{MC}, \mathfrak{t})$ at $\mathbf{1}_{x_0}$ and the germ of (M_1, θ_1, h_1) at m_1 can be identified with the germ of $(\mathfrak{s}^{-1}(x_0), \omega_{MC}, \mathfrak{t})$ at g . We conclude that (M_0, θ_0, h_0) and (M_1, θ_1, h_1) are both germs of the realization $(\mathfrak{s}^{-1}(x_0), \omega_{MC}, \mathfrak{t})$.

Conversely, suppose that there exists a connected realization (M, θ, h) such that (M_0, θ_0, h_0) and (M_1, θ_1, h_1) are the germs of θ at points p_0 and p_1 of M , respectively. Let γ be a curve joining p_0 to p_1 and cover it by a finite family U_1, \dots, U_k of open sets of M with the property that the restriction of θ to each U_i is equivalent to the restriction of the Maurer-Cartan form to an open set of some \mathfrak{s} -fiber of \mathcal{G} , i.e., $\theta|_{U_i} = \phi_i^* \omega_{MC}$ for some diffeomorphism $\phi_i : U_i \rightarrow \phi_i(U_i) \subset \mathfrak{s}^{-1}(x_i)$.

We proceed by induction on the number of open sets needed to join p_0 to p_1 . Suppose that both p_0 and p_1 belong to the same open set U_1 . Then $\phi_1(p_0)$ and $\phi_1(p_1)$ are both on the same \mathfrak{s} -fiber. Thus, $h(p_0) = \mathfrak{t} \circ \phi_1(p_0)$ belongs to the same orbit as $h(p_1) = \mathfrak{t} \circ \phi_1(p_1)$. Now assume that the result is true for $k - 1$ open sets. Then any point q in U_{k-1} is mapped by h to the same orbit of $h(p_0)$. Let q be a point in $U_{k-1} \cap U_k$. Then, on one hand, since q and p_1 belong to U_k it follows that h maps them both to the same orbit of A . On the other hand, by the inductive hypothesis, it follows that h also maps p_0 and q to the same orbit of A . \square

Remark 6.2. *The proposition above may not be true when A is not weakly integrable. The problem is that in this case, the global object associated to A_L is only a topological groupoid which is smooth only in a neighborhood of the identity section. Thus, if $x, y \in X$ are points in the same orbit which are “too far away”, then we might not be able to find a differentiable realization “covering” both points at the same time.*

Motivated by this remark, it is natural to consider the problem of existence of realizations (M, θ, h) of a Cartan’s problem, such that the image of h is the whole leaf of the classifying Lie algebroid.

Definition 6.3. *A connected realization (M, θ, h) is called **full** if h is surjective onto the orbit of A that it “covers”.*

The paradigmatic example of a full realization is obtained as follows.

Example 6.4. *Assume that the classifying Lie algebroid A is weakly integrable. Then the \mathfrak{s} -fibers of the Weinstein groupoid $\mathcal{G}(A)$ are smooth manifolds and carry (the restriction of) the Maurer-Cartan form. Each triple $(\mathfrak{s}^{-1}(x), \omega_{MC}, \mathfrak{t})$ is a full realization.*

In fact, there is a partial converse to the previous example. To explain this we also need the following:

Definition 6.5. *A realization (M, θ, h) of a Cartan’s problem is said to be **complete** if it is a full realization and every local equivalence $\phi : U \rightarrow V$, defined on open sets $U, V \subset M$ can be extended to a global equivalence $\tilde{\phi} : M \rightarrow M$, such that $\tilde{\phi}|_U = \phi$.*

Example 6.6 ((Example 6.4 continued)). *By Corollary 4.9, every local equivalence of the realization $(\mathfrak{s}^{-1}(x), \omega_{MC}, \mathfrak{t})$ is the restriction of right translation by same element $g \in \mathcal{G}(A)$. Hence, this is an example of a complete realization.*

Now we can state the following characterization (an analogous situation occurs in [16]):

Proposition 6.7. *Let $A \rightarrow X$ be the classifying Lie algebroid of a Cartan’s realization problem, and let $L \subset X$ be an orbit of A . Then there exists a complete realization over L if and only if the restriction $A|_L$ is integrable.*

Proof. Assume that $A|_L$ is integrable by $\mathcal{G} \rightrightarrows L$. Then, as observed above, for any $x \in L$, the realization $(\mathfrak{s}^{-1}(x), \omega_{MC}, \mathfrak{t})$ is complete.

Conversely, let (M, θ, h) be a complete realization which covers L . By Theorem 5.8, the group of symmetries $G = \text{Diff}(M, \theta, h)$ is a Lie group of transformations of M which fixes the fibers of $h : M \rightarrow L$. If $x, y \in M$ are such that $h(x) = h(y)$ then, by Theorem 5.3, there exists a local equivalence $\phi : U \rightarrow V$ such that $\phi(x) = y$. By the completeness of the realization, ϕ can be extended to a global equivalence $\tilde{\phi} \in G$. We conclude that G acts transitively on the fibers of h .

Now we claim that the action of G on M is free. This will follow by showing that the fixed point set of a symmetry $\phi : M \rightarrow M$ is both open and closed. It

is obviously closed. In order to prove that it is also open, we consider the vector fields $\{X_i\}$ dual to the coframe $\{\theta^i\}$ and denote by $\phi_{X_i}^t$ their (local) flows. Then, since ϕ is a symmetry, we have:

$$\phi_* X_i = X_i \quad \Leftrightarrow \quad \phi \circ \phi_{X_i}^t = \phi_{X_i}^t \quad (\text{whenever defined}).$$

Therefore, if $p \in M$ is some fixed point of ϕ , it follows that $\phi_{X_i}^t(p)$ is also a fixed point of ϕ . Since the X_i are linearly independent everywhere, it follows that there is neighborhood of p consisting of fixed points. This shows that the fixed point set of ϕ is open, and the claim follows.

We conclude that the G -action on M is free and its orbits are the fibers of $h : M \rightarrow L$, so that we have a principal bundle

$$\begin{array}{c} M \xrightarrow{\quad} G \\ \downarrow h \\ L \end{array}$$

The Atiyah algebroid of this principal bundle is isomorphic to $A|_L$. Hence, $A|_L$ is integrable as claimed. \square

6.2. The Global Classification Problem. We now turn to the global classification of coframes (Problem 1.9). For the remainder of this section, (n, X, C_{ij}^k, F_i) denotes a realization problem with classifying Lie algebroid $A \rightarrow X$ and (M, θ, h) is a realization.

The global equivalence of realizations of a Cartan's problem is much more delicate than the question of local equivalence. The reason is that the classifying Lie algebroid will not distinguish between a realization and its covering. For example, Olver in [14] constructs a simply connected manifold with a Lie algebra valued Maurer-Cartan form which cannot be globally embedded into any Lie group, but which is locally equivalent at every point to an open set of a Lie group.

We have the following theorem (the special cases of coframes of rank zero or maximal rank were considered by Olver in [13]):


Theorem 6.8. *Let (M, θ, h) be a realization of a Cartan problem and suppose that the classifying Lie algebroid $A \rightarrow X$ is weakly integrable. Then M is globally equivalent up to cover to an open set of an \mathfrak{s} -fiber of a groupoid \mathcal{G} integrating A .*

Proof. Let $L \subset X$ be the orbit containing $h(M)$ and choose a Lie groupoid \mathcal{G}_L integrating $A|_L$. Also, let \mathcal{D} be the distribution on $M_{h \times \mathfrak{t}} \mathcal{G}_L$ considered in the proof of Theorem 4.6 and let N be a maximal integral manifold of \mathcal{D} . If $\pi_M : M_{h \times \mathfrak{t}} \mathcal{G}_L \rightarrow M$ and $\pi_{\mathcal{G}} : M_{h \times \mathfrak{t}} \mathcal{G}_L \rightarrow \mathcal{G}_L$ denote the natural projections, then $\pi_{\mathcal{G}}(N)$ is totally contained in a single \mathfrak{s} -fiber of \mathcal{G}_L , say, $\mathfrak{s}^{-1}(x)$. Moreover, the restrictions of the projections to N , $\pi_M : N \rightarrow M$ and $\pi_{\mathcal{G}} : N \rightarrow \mathfrak{s}^{-1}(x)$ are local diffeomorphisms.

We claim that N , equipped with the coframe $\pi_M^* \theta = \pi_{\mathcal{G}}^* \omega_{\text{MC}}$, is a common realization cover of M and an open set of an \mathfrak{s} -fiber of \mathcal{G}_L . To show this, all we must show is that $\pi_M(N) = M$. In fact, if this is true, M will be globally equivalent up to cover to $\pi_{\mathcal{G}}(N) \subset \mathfrak{s}^{-1}(x)$.

To prove this, suppose that $\pi_M(N)$ is a proper submanifold of M and let $p_0 \in M - \pi_M(N)$ be a point in the closure of $\pi_M(N)$ (remember that $\pi_M|_N$ is an open map). Then, by the local universal property of Maurer-Cartan forms, there is an open set U of p_0 in M and a diffeomorphism $\phi_0 : U \rightarrow \phi(U) \subset \mathfrak{s}^{-1}(h(p_0))$ such that $\phi_0^* \omega_{\text{MC}} = \theta$ and $\phi_0(p_0) = \mathbf{1}_{h(p_0)}$. It follows that the graph of ϕ_0 is also an integral manifold N_0 of the distribution \mathcal{D} on $M_{h \times \mathfrak{t}} \mathcal{G}$ which passes through $(p_0, \mathbf{1}_{h(p_0)})$. Now let $p = \pi_M(p, g)$ be any point in $U \cap \pi_M(N)$ where $(p, g) \in N$. Then $\phi_0(p) \in \mathcal{G}$ is an

arrow from $h(p_0)$ to $h(p)$ and g is an arrow from x to $h(p)$ and thus $g_0 = \phi_0(p)^{-1} \cdot g$ is an arrow from x to $h(p_0)$, i.e.,



Now, by virtue of the invariance of the Maurer-Cartan form, the manifold

$$R_{g_0}N_0 = \{(\bar{p}, \bar{g} \cdot g_0) : (\bar{p}, \bar{g}) \in N_0\}$$

is an integral manifold of \mathcal{D} . But then, the point $(p, g) = (p, \phi_0(p) \cdot g_0)$ belongs to $R_{g_0}N_0$ and to N , and thus, by the uniqueness and maximality of N it follows that N contains $R_{g_0}N_0$. But then, $(p_0, \phi_0(p_0) \cdot g_0)$ is a point of N which projects through π_M to p_0 , which contradicts the fact that $p_0 \in M - \pi_M(N)$, proving the theorem. \square

We can now deduce the special cases where the rank of the coframe is either zero or maximal:

Corollary 6.9 (Theorem 14.28 of [13]). *Let θ be a coframe of rank 0 on a manifold M . Then M is globally equivalent, up to covering, to an open set of a Lie group.*

Proof. If θ has rank 0, the classifying Lie algebroid of the coframe has base $X = \{*\}$, i.e., it is a Lie algebra, so it is integrable by a Lie group. It follows that M is globally equivalent, up to covering, to an open set of this Lie group. \square

Corollary 6.10 (Theorem 14.30 of [13]). *Let (M_1, θ_1, h_1) and (M_2, θ_2, h_2) be full realizations of rank n over the same leaf $L \subset X$ of the classifying Lie algebroid $A \rightarrow X$. Then M_1 and M_2 are realization covers of a common realization (M, θ, h) .*

Proof. The assumptions of the corollary guarantee that L is a n -dimensional leaf of A for which $h_1(M_1) = L = h_2(M_2)$. Since A has rank n , it follows that the anchor of the restriction of A to L is injective, and thus $A|_L$ is integrable (see [5]). Moreover, its Lie groupoid \mathcal{G} is étale, i.e., its source and target maps are local diffeomorphisms.

It follows that we can make $M = L$ into a realization by equipping it with the pullback of the Maurer-Cartan form by the local inverses of \mathbf{t} . More precisely, let U be an open set in $\mathbf{s}^{-1}(x)$ for which the restriction of \mathbf{t} is one-to-one and let $V = \mathbf{t}(U)$ be the open image of U by \mathbf{t} . Define θ_U to be the coframe on V given by

$$\theta_U = (\mathbf{t}|_U^{-1})^* \omega_{\text{MC}}.$$

Then θ_U is the restriction to V of a globally defined coframe on L . In fact, suppose that \bar{U} is another open set of $\mathbf{s}^{-1}(x)$ for which $\mathbf{t}|_{\bar{U}}$ is one-to-one and such that $\bar{V} = \mathbf{t}(\bar{U})$ intersects V . Denote by $\theta_{\bar{U}}$ the coframe on \bar{V} defined by

$$\theta_{\bar{U}} = (\mathbf{t}|_{\bar{U}}^{-1})^* \omega_{\text{MC}}.$$

We will show that θ_U and $\theta_{\bar{U}}$ coincide on the intersection $V \cap \bar{V}$. After shrinking U and \bar{U} , if necessary, we may assume that $V = \bar{V}$. But then, since \mathcal{G} is étale, it follows that the isotropy group \mathcal{G}_x is discrete, which implies that U is the right translation of \bar{U} by an element $g \in \mathcal{G}_x$, i.e., $U = R_g(\bar{U})$. Thus,

$$\begin{aligned} \theta_U &= (R_g \circ \mathbf{t}|_{\bar{U}}^{-1})^* \omega_{\text{MC}} \\ &= (\mathbf{t}|_{\bar{U}}^{-1})^* (R_g^* \omega_{\text{MC}}) \\ &= (\mathbf{t}|_{\bar{U}}^{-1})^* \omega_{\text{MC}} = \theta_{\bar{U}}, \end{aligned}$$

and θ is a well defined global coframe on L .

Now, since both M_1 and M_2 are globally equivalent, up to covering, to open sets in $\mathfrak{s}^{-1}(x)$, it follows that the surjective submersions $h_i : M_i \rightarrow L$ are realization covers. \square

Remark 6.11. *There is a different, more direct, argument to see that the h_i 's are realization covers. In fact, the coframe θ_1 on M_1 is locally the pullback of the Maurer-Cartan form on an \mathfrak{s} -fiber of \mathcal{G} by some locally defined diffeomorphism $\phi_1 : W_1 \subset M_1 \rightarrow \mathfrak{s}^{-1}(x)$. After shrinking W_1 , we may assume that the restriction of \mathfrak{t} to $U_1 = \phi_1(W_1)$ is a diffeomorphism. But then,*

$$\begin{aligned} h_1^* \theta &= (\mathfrak{t} \circ \phi_1)^* \theta \\ &= \phi_1^* (\mathfrak{t}^* \circ (\mathfrak{t}|_{U_1}^{-1})^* \omega_{MC}) \\ &= \phi_1^* \omega_{MC} = \theta_1. \end{aligned}$$

Obviously, the very same argument can be given to show that h_2 is also a realization cover. This can be summarized by the diagram:

$$\begin{array}{ccc} & \mathfrak{s}^{-1}(x) & \\ \phi_1 \nearrow & \downarrow \mathfrak{t} & \nwarrow \phi_2 \\ M_1 & & M_2 \\ \searrow h_1 & & \swarrow h_2 \\ & L & \end{array}$$

6.3. Cohomological Invariants of Geometric Structures. The classifying Lie algebroid A of a fixed geometric structure should be seen as a basic invariant of global equivalence up to covering of the structure. In fact, we have the following result:

Proposition 6.12. *Let θ be a fully regular coframe on M and let $\bar{\theta}$ be an arbitrary coframe on \bar{M} . If (M, θ) and $(\bar{M}, \bar{\theta})$ are globally equivalent, up to covering, then*

- (i) $\bar{\theta}$ is a fully regular coframe on \bar{M} , and
- (ii) the classifying Lie algebroids of θ and $\bar{\theta}$ are isomorphic.

Proof. In order to prove this proposition, it is enough to consider the case where θ is a fully regular coframe on M , $\bar{\theta}$ is a coframe on another manifold \bar{M} and $\pi : \bar{M} \rightarrow M$ is a surjective local diffeomorphism which preserves the coframes.

First we need:

Lemma 6.13. *The invariant functions of the two coframes are in 1:1 correspondence: $\text{Inv}(\bar{\theta}) = \pi^* \text{Inv}(\theta)$.*

Proof of Lemma 6.13. Clearly, since $\pi : \bar{M} \rightarrow M$ is a cover, the locally defined self equivalences of (M, θ) and $(\bar{M}, \bar{\theta})$ correspond to each other. Hence the result follows. \square

It follows that if θ is a fully regular coframe so is the coframe $\bar{\theta}$. Let us denote by $A_\theta \rightarrow X_\theta$ and $A_{\bar{\theta}} \rightarrow X_{\bar{\theta}}$ the classifying algebroids of (M, θ) and $(\bar{M}, \bar{\theta})$, respectively. Also, we denote by $\kappa_\theta : M \rightarrow X_\theta$ and $\kappa_{\bar{\theta}} : \bar{M} \rightarrow X_{\bar{\theta}}$ the corresponding classifying maps, so $(M, \theta, \kappa_\theta)$ and $(\bar{M}, \bar{\theta}, \kappa_{\bar{\theta}})$ are realizations of A_θ and $A_{\bar{\theta}}$. Since $(\bar{M}, \bar{\theta}, \kappa_{\bar{\theta}} \circ \pi)$ is also a realization of A_θ it follows from Corollary 5.11 that there is an algebroid morphism from A_θ to $A_{\bar{\theta}}$, which is injective on the fibers and covers

a local diffeomorphism $X_\theta \rightarrow X_{\bar{\theta}}$ making the following diagram commutative:

$$\begin{array}{ccc} M & \xleftarrow{\pi} & \bar{M} \\ \kappa_\theta \downarrow & & \downarrow \kappa_{\bar{\theta}} \\ X_\theta & \longrightarrow & X_{\bar{\theta}} \end{array}$$

We claim that the map $X_\theta \rightarrow X_{\bar{\theta}}$ is injective, so that A_θ and $A_{\bar{\theta}}$ are isomorphic. In fact, let $x, y \in X_\theta$ which are mapped to the same point in $z \in X_{\bar{\theta}}$. Then we can choose points $p, q \in M$ and $\bar{p}, \bar{q} \in \bar{M}$ which are mapped to each other under the commutative diagram above. Then, by Theorem 5.3, there exists a local equivalence $\bar{\phi} : \bar{M} \rightarrow \bar{M}$, such that $\bar{\phi}(\bar{p}) = \bar{q}$. But then (after restricting to small enough open sets) $\bar{\phi}$ covers a local equivalence $\phi : M \rightarrow M$ such that $\phi(p) = q$. This means that p and q are local equivalent, so $x = \kappa_\theta(p) = \kappa_\theta(q) = y$, thus proving the claim. \square

Even though the isomorphism class of the classifying Lie algebroid does not distinguish coframes which are globally equivalent, up to covering, one can use its Lie algebroid cohomology to obtain invariants of coframes. To illustrate this point of view, we will now describe two invariants of coframes arising in this way.

Given a Lie algebroid A we will denote by $(\Omega^\bullet(A), d_A)$ the complex of A -forms where $\Omega^\bullet(A) := \Gamma(\wedge^\bullet A^*)$ and the differential is given by the formula:

$$\begin{aligned} d_A \omega(\alpha_0, \dots, \alpha_k) &= \sum_{i=1}^k (-1)^{i\sharp}(\alpha_i) \cdot \omega(\alpha_0, \dots, \hat{\alpha}_i, \dots, \alpha_k) + \\ &+ \sum_{0 \leq i < j \leq k} (-1)^{i+j} \omega([\alpha_i, \alpha_j], \alpha_0, \dots, \hat{\alpha}_i, \dots, \hat{\alpha}_j, \dots, \alpha_k). \end{aligned}$$

The resulting cohomology $H^\bullet(A)$ is called the Lie algebroid cohomology of A . Every Lie algebroid morphism $\Phi : A \rightarrow B$ induces a chain map

$$\Phi^* : (\Omega^\bullet(B), d_B) \rightarrow (\Omega^\bullet(A), d_A),$$

and, hence, also a map in cohomology $\Phi^* : H^\bullet(B) \rightarrow H^\bullet(A)$.

For a fully regular coframe θ on M the Lie algebroid cohomology of its classifying algebroid A_θ can be described in terms of *invariant forms*. Let us call $\omega \in \Omega^k(M)$ a **k-invariant form** if for any locally defined equivalence of the coframe $\phi : M \rightarrow M$ one has:

$$\phi^* \omega = \omega.$$

Since the differential of an invariant form is again an invariant form, the subspace of invariant forms $\Omega_\theta^*(M)$ is a subcomplex of the de Rham complex.

Definition 6.14. *The invariant cohomology of (M, θ) , denoted $H_\theta^*(M)$ is the cohomology of the complex of invariant forms.*

Now we note that:

Proposition 6.15. *Let (M, θ) be a fully regular coframe. Then the Lie algebroid cohomology $H^*(A_\theta)$ coincides with the invariant cohomology $H_\theta^*(M)$.*

Proof. This is simply a consequence of the fact that $\theta^* : \Omega^\bullet(A_\theta) \rightarrow \Omega^\bullet(M)$ is injective and has image precisely the complex of invariant forms. In fact, a form $\omega \in \Omega^k(M)$ is the pullback by θ of a section φ of $\wedge^k A_\theta^*$ iff when we write ω in terms of the coframe θ we have:

$$\omega = \sum_{i_1 < \dots < i_k} a_{i_1, \dots, i_k} \circ \kappa_\theta \theta^{i_1} \wedge \dots \wedge \theta^{i_k},$$

for some smooth functions $a_{i_1, \dots, i_k} \in C^\infty(X_\theta)$. \square

Since can view the coframe as a Lie algebroid map $\theta : TM \rightarrow A_\theta$ (over the classifying map $\kappa_\theta : M \rightarrow X_\theta$), there is an induced map in cohomology $\theta^* : H^\bullet(A_\theta) \rightarrow H_{\text{dR}}^\bullet(M)$. By the previous proposition, a class belongs to the image of this map iff it can be represented by an invariant k -form. In fact, under the isomorphism $H^*(A_\theta) \simeq H_\theta^*(M)$ the map $\theta^* : H^\bullet(A_\theta) \rightarrow H_{\text{dR}}^\bullet(M)$ corresponds to the map $H_\theta^*(M) \rightarrow H_{\text{dR}}^\bullet(M)$ induced by the forgetful map.

Remark 6.16. *If every local equivalence can be extended to a global equivalence (i.e., if $(M, \theta, \kappa_\theta)$ is a complete realization) then $H_\theta^*(M)$ coincides with the invariant cohomology $H_G^\bullet(M)$ associated with the action of $G = \text{Diff}(M, \theta)$, the group of symmetries of the coframe. If further G is compact, then we have $H_G^\bullet(M) = H_{\text{dR}}^\bullet(M)$.*

From Proposition 6.12 we see that invariant cohomology is an invariant of global equivalence, up to covering.

Corollary 6.17. *Let (M, θ) and $(\bar{M}, \bar{\theta})$ be fully regular coframes which are globally equivalent, up to covering. Then their invariant cohomologies are isomorphic $H_\theta^*(M) \simeq H_{\bar{\theta}}^*(\bar{M})$.*

The invariant cohomology of a coframe is the natural place where classes associated with the coframe live. We will study next one such example.

Remark 6.18. *There is also an obvious extension of these constructions and results to any realization (M, θ, h) of a Cartan problem, so one can define the invariant cohomology $H_{\theta, h}^*(M)$. This cohomology is invariant under global equivalence and if (M, θ, h) is complete, and $G = \text{Diff}(M, \theta, h)$, then it coincides with $H_G^\bullet(M)$.*

6.4. The modular class. Lie algebroids have associated intrinsic characteristic classes (see [9, 4, 6]). Therefore, for a fully regular coframe θ we can consider the characteristic classes of its classifying Lie algebroid A_θ and we obtain certain invariant cohomology classes which maybe called the characteristic classes of the coframe. The simplest of these classes is the *modular class* ([8, 12]) which we recall briefly.

Given a Lie algebroid $A \rightarrow M$ the line bundle $L_A := \wedge^{\text{top}} A \otimes \wedge^{\text{top}} T^*M$ carries a natural flat A -connection, which makes L into a A -representation, defined by:

$$\nabla_\alpha(\omega \otimes \nu) = \mathcal{L}_\alpha \omega \otimes \nu + \omega \otimes \mathcal{L}_{\rho(\alpha)} \nu \quad (\alpha \in \Gamma(A)).$$

When L_A is orientable, so that it carries a nowhere vanishing section $\mu \in \Gamma(L_A)$, we have:

$$\nabla_\alpha \mu = \langle c_\mu, \alpha \rangle \mu, \quad \forall \alpha \in \Gamma(A),$$

for a 1-form $c_\mu \in \Omega^1(A)$. One checks easily that c_μ is d_A -closed and one calls c_μ the **modular cocycle** of A relative to the nowhere vanishing section μ . If $\mu' = f\mu$ is some other non-vanishing section, then one finds that:

$$c_{\mu'} = c_\mu + d_A \log |f|.$$

so the cohomology class $\text{mod}(A) := [c_\mu] \in H^1(A)$ does not depend on the choice of global section μ . One calls $\text{mod}(A)$ the **modular class** of A . When L_A is not orientable, one repeats the construction for $L_A \otimes L_A$ and defines $\text{mod}(A)$ to be one half the cohomology class associated with the line bundle $L_A \otimes L_A$.

Definition 6.19. *The modular class of a fully regular coframe is the invariant cohomology class $\text{mod}(\theta) \in H_\theta^1(M)$ which under the natural isomorphism $H_\theta^*(M) \simeq H^\bullet(A_\theta)$ corresponds to the class $\text{mod}(A_\theta)$.*

It should be clear that if (M, θ) and $(\bar{M}, \bar{\theta})$ are fully regular coframes which are globally equivalent, up to covering, then under the natural isomorphism $H_\theta^*(M) \simeq H_{\bar{\theta}}^*(\bar{M})$ given by Corollary 6.17 the modular classes correspond to each other.

Our next proposition leads to a geometric interpretation of the modular class of a coframe:

Proposition 6.20. *Let (M, θ) be a fully regular coframe with symmetry Lie algebra $\mathfrak{g} := \mathfrak{X}(M, \theta)$ of dimension k . Let $\{\xi_1, \dots, \xi_k\}$ be a basis of \mathfrak{g} and let μ_G be an invariant k -form which restricts to a volume form on each \mathfrak{g} -orbit. Then:*

$$\text{mod}(\theta) = -[\text{d log } |\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle|] \in H_\theta^1(M).$$

Proof. For a transitive algebroid the bundle $\ker \rho$ is a natural A -representation and we have a natural isomorphism of A -representations $\wedge^{\text{top}} \ker \rho \simeq L_A$. Observe also that the classifying map $k_\theta : M \rightarrow X$, whose fibers coincide with the \mathfrak{g} -orbits, give an identification $k_\theta^* \ker \rho \simeq \ker dk_\rho$. Hence, we can identify invariant sections of $\ker dk_\rho$ with sections of $\ker \rho$, and it follows that any invariant k -form μ_G which restricts to a volume form on each \mathfrak{g} -orbit determines a global section of L_A^* (and, hence, of L_A). If $\{\alpha_1, \dots, \alpha_n\}$ denotes the canonical basis of A (which we identify with the invariant vector fields dual to the coframe $\{\theta^1, \dots, \theta^n\}$), then under these identifications, we have that:

$$\nabla_{\alpha_i} \mu_G = \mathcal{L}_{\alpha_i} \mu_G,$$

where on the right-hand side we view α_i as an invariant vector field on M . It follows that the modular cocycle relative to the section $\mu = \mu_G^*$ is given by:

$$\begin{aligned} c_\mu(\alpha_i) &= \frac{\langle c_\mu(\alpha_i) \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle}{\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle} \\ &= \frac{\langle -\nabla_{\alpha_i} \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle}{\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle} \\ &= \frac{\langle -\mathcal{L}_{\alpha_i} \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle}{\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle} \\ &= -\frac{\mathcal{L}_{\alpha_i} \langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle}{\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle} = -\langle \alpha_i, \text{d log } |\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle| \rangle \end{aligned}$$

This proves that the expression for the modular class in the statement of the proposition holds. \square

The proposition shows, in particular, that under the natural map $H_\theta^*(M) \rightarrow H_{\text{dR}}^\bullet(M)$ the modular class $\text{mod}(\theta)$ is mapped to zero. Thus, the modular class is an obstruction for a complete coframe to have a compact symmetry group, i.e.,

Corollary 6.21. *Let (M, θ) be a fully regular coframe. If θ is complete, and the symmetry group $\text{Diff}(M, \theta)$ is compact, then $\text{mod}(\theta) = 0$.*

Also, we conclude from Proposition 6.20 that:

Corollary 6.22. *The symmetry Lie algebra $\mathfrak{g} := \mathfrak{X}(M, \theta)$ is unimodular if and only if $\text{mod}(\theta) = 0$.*

Proof. One finds for any $\xi \in \mathfrak{g}$ that:

$$\begin{aligned} \mathcal{L}_\xi \log |\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle| &= \frac{\mathcal{L}_\xi \langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle}{\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle} \\ &= \frac{\sum_{i=1}^k \langle \mu_G, \xi_1 \wedge \dots \wedge [\xi, \xi_i] \wedge \dots \wedge \xi_k \rangle}{\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle} = \text{trace}(\text{ad } \xi) \end{aligned}$$

Hence, we conclude that $\log |\langle \mu_G, \xi_1 \wedge \dots \wedge \xi_k \rangle|$ is an invariant function if and only if \mathfrak{g} is unimodular. \square

Remark 6.23. *More generally, if (M, θ, h) is a realization of a Cartan problem associated with a classifying Lie algebroid $A \rightarrow X$ one can also define its modular class by*

$$\text{mod}(M, \theta, h) := \text{mod}(A) \in H_{\theta, h}^1(M)$$

where $H_{\theta, h}^1(M)$ denotes the invariant cohomology (see Remark 6.18). A version of Proposition 6.20 also holds.

7. AN EXAMPLE: SURFACES OF REVOLUTION

In this section, we consider the local classification of surfaces of revolutions up to isometries. We begin by deducing the structure equations of a generic surface of revolution following very closely the differential analysis presented in Chapter 12 of [13]. However, since the local moduli space of surfaces of revolution is infinite dimensional (we will see that it depends on an arbitrary function which will be denoted by H), we will then restrict to a finite dimensional class of surfaces of revolution, namely those for which the moduli H has a fixed value, and exhibit its classifying Lie algebroid. We use this example mainly to illustrate several of the concepts and results presented throughout the paper. In the sequel to this paper, where we discuss G -structures, we will discuss more sophisticated (and interesting examples), such as special symplectic manifolds.

Let x, y, z denote the canonical coordinates on \mathbb{R}^3 and let $z = f(x)$ be a curve in the plane $\{y = 0\}$ which does not intersect the z -axis, i.e., such that $f(x) \neq 0$ for all $x \in \mathbb{R}$. Let Σ be the surface obtained by rotating $f(x)$ around the z -axis, which we can parameterize using polar coordinates

$$\begin{cases} x = r \cos v \\ y = r \sin v \\ z = f(r) \end{cases}$$

where $r > 0$.

The relevant coframes for this classification problem is the one which diagonalizes the metric ds^2 induced by the Euclidean metric on $\Sigma \subset \mathbb{R}^3$, i.e., the coframes $\{\omega^1, \omega^2\}$ for which

$$ds^2 = (\omega^1)^2 + (\omega^2)^2.$$

In order to simplify the expression for such a coframe, we first perform the change of coordinates

$$u = \int \sqrt{1 + f'(r)^2} dr$$

with inverse $r = h(u)$, where $h(u) > 0$ for all u . With respect to the new coordinates (u, v) , the metric can be written as

$$ds^2 = (du)^2 + h(u)^2 (dv)^2,$$

so any diagonalizing coframe for ds^2 is given by

$$\begin{cases} \omega^1 = du \\ \omega^2 = h(u)dv. \end{cases}$$

Notice that any coframe which differs from $\{\omega^1, \omega^2\}$ by a rotation represents the same metric. This suggests that we should consider the coframe

$$\begin{cases} \theta^1 = (\sin t)\omega^1 + (\cos t)\omega^2 \\ \theta^2 = (-\cos t)\omega^1 + (\sin t)\omega^2 \\ \alpha = dt \end{cases}$$

on $\Sigma \times S^1$, where α is the Maurer-Cartan form on $S^1 \cong \text{SO}(2)$.

Remark 7.1. *The passage from M to $M \times G$ is the very first preliminary step in Cartan's equivalence method. The coframe in $M \times G$, obtained by adding the Maurer-Cartan forms, is called the lifted coframe (see [13]).*

Next, in order to obtain the structure equations and structure functions, we differentiate the one forms θ^1 , θ^2 and α obtaining

$$(7.1) \quad \begin{cases} d\theta^1 = -\alpha \wedge \theta^2 + \left(\frac{h'(u)}{h(u)} \cos t\right) \theta^1 \wedge \theta^2 \\ d\theta^2 = \alpha \wedge \theta^1 + \left(\frac{h'(u)}{h(u)} \sin t\right) \theta^1 \wedge \theta^2 \\ d\alpha = 0 \end{cases}$$

These structure equations depend explicitly on the coordinates u and t . However, this can be fixed by replacing the form α by

$$(7.2) \quad \eta = \alpha - \frac{h'(u)}{h(u)} ((\cos t)\theta^1 + (\sin t)\theta^2) = \alpha - h'(u)dv.$$

If we now rewrite (7.1) in terms of the new coframe we obtain

$$\begin{cases} d\theta^1 = -\eta \wedge \theta^2 \\ d\theta^2 = \eta \wedge \theta^1 \\ d\eta = \kappa \theta^1 \wedge \theta^2 \end{cases}$$

where $\kappa = -h''(u)/h(u)$ is the Gaussian curvature of Σ .

Remark 7.2. *The substitution of α by η is another step in Cartan's equivalence method and is known as absorption of torsion (see [13]).*

The next step is to calculate the coframe derivative of κ . We obtain

$$d\kappa = \kappa'(u)((\sin t)\theta^1 - (\cos t)\theta^2).$$

However, since Σ is a *surface of revolution*, we know that its symmetry Lie group must be at least one dimensional. It follows that there can be at most two independent invariants, and since we already have two invariant functions, namely t and κ , we conclude that $\kappa'(u) = H(\kappa)$ for some $H \in C^\infty(\mathbb{R})$.

Remark 7.3. *By its definition, t is the angle between the orthogonal coframe at a point and the special coframe $\{du, h(u)dv\}$ given in the adapted coordinates. Thus, imposing t as an invariant is the same as considering surfaces of revolution up to isometries which fix the angle t . This can be geometrically interpreted as a restriction on the allowed symmetries of Σ . For a generic surface of revolution this is not really a restriction but, for example, when $H = 0$, i.e., when the curvature of Σ is constant, this imposes a true restriction on the symmetry group.*

Finally, from (7.2) we find that

$$dt = \eta + J(\kappa)((\cos t)\theta^1 + (\sin t)\theta^2)$$

for some $J \in C^\infty(\mathbb{R})$. The relationship between the two functions H and J follows from imposing $d^2\kappa = 0$. If we assume $\kappa'(u) \neq 0$, i.e., that $H(\kappa) \neq 0$, it yields

$$(7.3) \quad J'(\kappa)H(\kappa) = -\kappa - J(\kappa)^2.$$

In conclusion, the full set of **structure equations for surfaces of revolution** is

$$(7.4) \quad \begin{cases} d\theta^1 = -\eta \wedge \theta^2 \\ d\theta^2 = \eta \wedge \theta^1 \\ d\eta = \kappa \theta^1 \wedge \theta^2 \\ d\kappa = H(\kappa)((\sin t)\theta^1 - (\cos t)\theta^2) \\ dt = \eta + J(\kappa)((\cos t)\theta^1 + (\sin t)\theta^2) \end{cases}$$

where H is an arbitrary smooth function and $J \in C^\infty(\mathbb{R})$ is determined by (7.3). Notice that $d^2 = 0$ is a consequence of the equations above.

Remark 7.4 (Continuing Remark 7.3). *When $\kappa'(u) = 0$ the analysis above collapses and the structure equations for surfaces of revolution reduce to:*

$$(7.5) \quad \begin{cases} d\theta^1 = -\eta \wedge \theta^2 \\ d\theta^2 = \eta \wedge \theta^1 \\ d\eta = \kappa\theta^1 \wedge \theta^2 \\ d\kappa = 0. \end{cases}$$

Now that we have found the structure equations for an arbitrary surface of revolution, we may immediately write the classifying Lie algebroid A for such surfaces. As a vector bundle, $A \cong (\mathbb{R} \times S^1) \times \mathbb{R}^3 \rightarrow \mathbb{R} \times S^1$. Its Lie bracket is given on the constant sections e_1, e_2 and e_3 by

$$\begin{aligned} [e_1, e_2](\kappa, t) &= -\kappa e_3 \\ [e_1, e_3](\kappa, t) &= e_2 \\ [e_2, e_3](\kappa, t) &= -e_1 \end{aligned}$$

and its anchor is given by

$$\begin{aligned} \sharp(e_1)(\kappa, t) &= H(\kappa) \sin t \frac{\partial}{\partial \kappa} + J(\kappa) \cos t \frac{\partial}{\partial t} \\ \sharp(e_2)(\kappa, t) &= -H(\kappa) \cos t \frac{\partial}{\partial \kappa} + J(\kappa) \sin t \frac{\partial}{\partial t} \\ \sharp(e_3)(\kappa, t) &= \frac{\partial}{\partial t}. \end{aligned}$$

Again, when $\kappa'(u) = 0$ (cf. Remarks 7.3 and 7.4), the analysis collapses and we obtain an algebroid over a 1-dimensional manifold with zero anchor (i.e., a bundle of Lie algebras).

For any function $H(\kappa)$ the equations above determine a Lie algebroid. From Theorem 5.3 and the fact that the algebroid is transitive, we deduce that:

Proposition 7.5. *For any $\kappa_0 \in \mathbb{R}$ and any $H \in C^\infty(\mathbb{R})$ there exists a surface of revolution Σ with $p \in \Sigma$ such that $\kappa(p) = \kappa_0$ and for which $\kappa'(u) = H(\kappa)$. Moreover, any two such surfaces of revolution are locally isometric in a neighborhood of the points corresponding to κ_0 .*

We can also use the classifying Lie algebroid to describe the infinitesimal symmetries of surfaces of revolution (see Proposition 5.7 and Theorem 5.8). For this, we must distinguish the cases $\kappa'(u) \neq 0$ and $\kappa'(u) = 0$:

$H(\kappa) \neq 0$:: In this case, which is generic, it is easy to see that the anchor is surjective, and thus the symmetry Lie algebra is 1-dimensional, which corresponds to rotation around the axis of revolution.

$H(\kappa) = 0$:: In this case, the surface of revolution has constant curvature. The orbits of the corresponding Lie algebroid are 0-dimensional, and thus, the symmetry Lie algebra is 3-dimensional. We have the following 3 possibilities:

\mathfrak{sl}_2	if $\kappa < 0$	Hyperbolic Geometry
\mathfrak{se}_2	if $\kappa = 0$	Euclidean Geometry
\mathfrak{so}_3	if $\kappa > 0$	Spherical Geometry

Remark 7.6. *Note that a surface of revolution which has non-constant $H(\kappa)$ which passes through 0 cannot appear as a realization of our Cartan's problem. The reason is that the corresponding coframe will not be fully regular.*

Remark 7.7. *Strictly speaking the classifying Lie algebroid described above (and also the conclusions deduced from it) concern the coframe θ^1, θ^2, η , and not the surface itself. However, it will be shown in [10] how to deal with Cartan's realization problem on finite type G -structures. In particular, the realization problem discussed here should be treated as a realization problem on the orthogonal frame bundle of the surface of revolution.*

We remark that whenever one fixes the function H , one obtains the classifying Lie algebroid discussed above is the classifying Lie algebroid of a single coframe; namely the coframe θ^1, θ^2, η on the orthogonal frame bundle of the surface of revolution. However, the (local) moduli space of surfaces of revolution is clearly infinite dimensional. Thus, in order to describe a class of surfaces of revolution, we will consider here only those for which H is constant.

Remark 7.8. *The condition that H is constant can be seen a second order differential equation on the curvature tensor R of the induced metric on the surface of revolution. In fact, using the coordinates u, v introduced earlier, we can express the class of surfaces being considered as the class of surfaces of revolution for which*

$$\nabla_{\frac{\partial}{\partial u}}(\nabla R) = 0,$$

where ∇ is the Levi-Civita connection of the surface. Since this condition is equivalent to $\kappa(u)$ being an affine map, we will denote such surfaces by **affinely curved surfaces of revolution**.

Note now that since we are not prescribing a specific value for the constant H , we must consider it as a new invariant function for affinely curved surfaces of revolution which is subject to the condition $dH = 0$. Similarly, J is no longer determined and must also be added to our set of invariant functions. It then follows from (7.3) that whenever $H \neq 0$ (which we assume from now on)

$$dJ = -(\kappa + J^2)((\sin t)\theta^1 - (\cos t)\theta^2).$$

Thus, as a complete set of structure equations for affinely curved surfaces of revolution one obtains:

$$(7.6) \quad \begin{cases} d\theta^1 = -\eta \wedge \theta^2 \\ d\theta^2 = \eta \wedge \theta^1 \\ d\eta = \kappa\theta^1 \wedge \theta^2 \\ d\kappa = H(\kappa)((\sin t)\theta^1 - (\cos t)\theta^2) \\ dt = \eta + J(\kappa)((\cos t)\theta^1 + (\sin t)\theta^2) \\ dH = 0 \\ dJ = -(\kappa + J^2)((\sin t)\theta^1 - (\cos t)\theta^2). \end{cases}$$

Again, it is easily verified that these structure equations have as a formal consequence that $d^2 = 0$, and thus they determine the classifying Lie algebroid B for affinely curved surfaces of revolution. As a vector bundle, $B \cong Y \times \mathbb{R}^3 \rightarrow Y$, where

$$Y = \{(\kappa, H, J, t) \in \mathbb{R}^3 \times S^1 : H \neq 0\}.$$

Its Lie bracket is given on the constant sections e_1, e_2 and e_3 by

$$\begin{aligned} [e_1, e_2](\kappa, H, J, t) &= -\kappa e_3 \\ [e_1, e_3](\kappa, H, J, t) &= e_2 \\ [e_2, e_3](\kappa, H, J, t) &= -e_1 \end{aligned}$$

and its anchor is given by

$$\begin{aligned} \sharp(e_1)(\kappa, H, J, t) &= H \sin t \frac{\partial}{\partial \kappa} - (\kappa + J^2) \sin t \frac{\partial}{\partial J} + J(\kappa) \cos t \frac{\partial}{\partial t} \\ \sharp(e_2)(\kappa, H, J, t) &= -H \cos t \frac{\partial}{\partial \kappa} + (\kappa + J^2) \cos t \frac{\partial}{\partial J} + J(\kappa) \sin t \frac{\partial}{\partial t} \\ \sharp(e_3)(\kappa, H, J, t) &= \frac{\partial}{\partial t}. \end{aligned}$$

Observe that the orbits of this Lie algebroid are 2-dimensional, so we can conclude that:

Proposition 7.9. *For any κ_0, H_0, J_0 with $H_0 \neq 0$ there exists an affinely curved surface of revolution whose structure invariants take the values κ_0, H_0, J_0 . Moreover,*

- (1) *Every affinely curved surface of revolution has a one dimensional symmetry Lie algebra.*
- (2) *If two affinely curved surfaces of revolution are open submanifolds of a third affinely curved surface of revolution, then there invariant constant function H must agree.*

APPENDIX A. THE CLASSIFYING LIE ALGEBROID OF A SINGLE COFRAME

In this appendix we will use the jet bundle approach to give a coordinate free construction of the classifying Lie algebroid associated to a fully regular coframe θ on a manifold M . We begin by recalling some general constructions.

Let N be a manifold and let $\pi : N \rightarrow M$ be a smooth map. Let us denote by $\text{Hom}(TN, \pi^*TM)$ the vector bundle over N whose sections are bundle maps

$$\begin{array}{ccc} TN & \longrightarrow & \pi^*TM \\ & \searrow & \swarrow \\ & N & \end{array}$$

and by $J^1(\text{Hom}(TN, \pi^*TM))$ the bundle of 1-jets of sections of $\text{Hom}(TN, \pi^*TM)$. We can identify a point of $J^1(\text{Hom}(TN, \pi^*TM))$ in the fiber over $x \in N$ with a pair (q, l_q) , where $q \in \text{Hom}_x(TN, \pi^*TM)$ and

$$l_q : T_x N \rightarrow \text{Hom}_x(TN, \pi^*TM)$$

is a linear map. It follows that we may define a bundle map

$$\begin{array}{ccc} J^1(\text{Hom}(TN, \pi^*TM)) & \xrightarrow{\text{Alt}} & \text{Hom}(\wedge^2 TN, \pi^*TM) \\ & \searrow & \swarrow \\ & N & \end{array}$$

by

$$\text{Alt}(q, l_q)(v \wedge w) = \frac{1}{2}l_q(v)(w) - l_q(w)(v).$$

Example A.1. *If we take M to be the real line \mathbb{R} and π to be a constant map, then the construction above yields a map*

$$\begin{array}{ccc} J^1(T^*N) & \xrightarrow{\text{Alt}} & \wedge^2 T^*N \\ & \searrow & \swarrow \\ & N & \end{array}$$

that satisfies

$$\text{Alt}(j^1\alpha) = d\alpha.$$

Now, note that we may view π_* as a section of $\text{Hom}(TN, \pi^*TM)$. Thus, by applying Alt , we obtain a section $\text{Alt}(j^1\pi_*) \in \text{Hom}(\wedge^2 TN, \pi^*TM)$.

Definition A.2. *The **structure tensor** of $\pi : N \rightarrow M$ is the map*

$$c : J^1N \rightarrow \text{Hom}(\wedge^2 TM, \pi^*TM)$$

defined by

$$c(H_p)(v \wedge w) = \text{Alt}(j^1\pi_*)(\tilde{v} \wedge \tilde{w}),$$

where \tilde{v} and \tilde{w} are the horizontal lifts of $v, w \in T_{\pi(p)}M$ to H_p .

Remark A.3. In the definition above, we have identified $j_x^1 s \in J^1 N$ with the horizontal subspace $H_{s(x)} = d_x s(T_x M)$ of $T_{s(x)} N$.

Example A.4. In the case where N is the frame bundle of M (or more generally a G -structure over M), the structure tensor is simply its first order structure function (see, for example, [19, 18] or [10]).

After these preliminary remarks, we now return to intrinsic construction of the classifying Lie algebroid of a fully regular coframe θ on a manifold M . Let us denote by $F(M)$ the frame bundle of M . Thus,

$$\begin{array}{ccc} F(M) & \xrightarrow{\quad} & \text{GL}_n \\ \pi \downarrow & & \\ M & & \end{array}$$

is a principal GL_n -bundle whose fiber over a point p is

$$\pi^{-1}(p) = \{ \phi : \mathbb{R}^n \rightarrow T_p M : \phi \text{ is a linear isomorphism} \}.$$

The coframe θ can be identified with a section of $F(M)$. It gives rise to trivializations $TM \cong M \times \mathbb{R}^n$ and $T^*M \cong M \times \mathbb{R}^n$ and with respect to the canonical base of \mathbb{R}^n , we can write $\theta = (\theta^1, \dots, \theta^n)$ or dually $\frac{\partial}{\partial \theta} = (\theta_1, \dots, \theta_n)$. A basis of sections of $\wedge^2 T^*M \otimes TM$ is then given by

$$\theta_k^{ij} = (\theta^i \wedge \theta^j) \otimes \theta_k.$$

With respect to this basis, the structure tensor c applied to $j^1 \theta$ gives rise to a section of $\wedge^2 T^*M \otimes TM$ whose coordinates are the structure functions, i.e.,

$$c(j^1 \theta) = \sum C_{ij}^k \theta_k^{ij}.$$

It follows that we may view the set $\mathcal{F}_0(M)$ of structure functions of θ as the section $c(j^1 \theta)$ of $\text{Hom}(\wedge^2 TM, TM)$.

Note also that $j^1 c$ is a map

$$J^1(J^1 F(M)) \rightarrow J^1(\text{Hom}(\wedge^2 TM, \pi^* TM)).$$

Recall that $J^1(\text{Hom}(\wedge^2 TM, \pi^* TM))$ is isomorphic to

$$(\wedge^2 T^*M \otimes TM) \times \text{Hom}(TM, \wedge^2 T^*M \otimes TM).$$

Let us denote by ∇ the flat connection on $\wedge^2 T^*M \otimes TM$ induced by θ . Then a basis of sections of $\text{Hom}(TM, \wedge^2 T^*M \otimes TM)$ is given by

$$\theta_{k,l}^{ij} = \nabla_{\theta_l} \theta_k^{ij}.$$

With respect to this basis, the section $j^1 c(j^2 \theta)$ of $J^1(\text{Hom}(\wedge^2 TM, TM))$ is written as

$$j^1 c(j^2 \theta) = \sum C_{ij}^k \theta_k^{ij} + \frac{\partial C_{ij}^k}{\partial \theta_l} \theta_{k,l}^{ij}.$$

Thus we may view the set $\mathcal{F}_1(M)$ formed by the structure functions and its coframe derivatives as the section $j^1 c(j^2 \theta)$ of

$$J^1(\text{Hom}(\wedge^2 TM, TM)).$$

If we continue in this way, we may identify the set $\mathcal{F}_r(M)$ of all coframe derivatives of the the structure functions up to order r with the section

$$j^{r-1} c(j^r \theta) \in \Gamma(J^r(\text{Hom}(\wedge^2 TM, TM))),$$

which we shall call the **r -th order structure section** of θ .

Now, the coframe θ induces a trivialization of $J^r(\text{Hom}(\wedge^2 TM, TM))$. Thus we have an identification between all its fibers. We shall denote them by \mathbb{K}_r and by

$$p_\theta^r : J^r(\text{Hom}(\wedge^2 TM, TM)) \rightarrow \mathbb{K}_r$$

the associated projection. We observe that the coframe θ is fully regular if and only if each of the maps

$$\begin{array}{ccc} M & \xrightarrow{j^{r-1}c(j^r\theta)} & \Gamma(J^r(\text{Hom}(\wedge^2 TM, TM))) \xrightarrow{p_\theta^r} \mathbb{K}_r \\ & \searrow & \nearrow \\ & & \mathbb{K}_r \end{array}$$

$\tilde{\kappa}_r$

has constant rank.

Thus, we can summarize the construction made so far by saying that each coframe θ on M , viewed as a section of $F(M)$, gives rise to a section

$$s_\theta \in \Gamma(J^\infty(\text{Hom}(\wedge^2 TM, TM))).$$

Moreover, since the coframe θ induces a trivialization of $J^\infty(\text{Hom}(\wedge^2 TM, TM))$, there is a natural projection

$$p_\theta : J^\infty(\text{Hom}(\wedge^2 TM, TM)) \rightarrow \mathbb{K}_\infty,$$

where \mathbb{K}_∞ denotes the fiber of $J^\infty(\text{Hom}(\wedge^2 TM, TM))$. If, additionally, we assume that θ is fully regular, then $\tilde{\kappa}_\theta = p_\theta \circ s_\theta$ has constant rank, and thus its image in \mathbb{K}_∞ is an immersed submanifold (possibly with self intersection) which we denote by \tilde{X} , i.e., there is a manifold X , and an immersion (not necessarily injective) $\phi : X \rightarrow \mathbb{K}_\infty$ such that

$$\begin{array}{ccc} M & \xrightarrow{s_\theta} & \Gamma(J^\infty(\text{Hom}(\wedge^2 TM, TM))) \xrightarrow{p_\theta} \mathbb{K}_\infty \\ & \searrow \tilde{\kappa}_\theta & \nearrow \phi \\ & & X \end{array}$$

$\tilde{\kappa}_\infty$

We remark that again, as in Section 2.2, we may take X to be the quotient of M by the equivalence relation where two points p and q are identified if and only if there is a locally defined formal equivalence of θ which takes p to q . Thus, by construction, we may view the structure section of order zero $c(j^1\theta) \in \Gamma(\text{Hom}(\wedge^2 TM, TM))$ as being defined on X . In other words, there is a smooth map

$$\hat{c} : X \rightarrow \text{Hom}(\wedge^2 TM, TM),$$

such that

$$\begin{array}{ccc} M & \xrightarrow{\tilde{\kappa}_\theta} & X \\ & \searrow j^1c(\theta) & \swarrow \hat{c} \\ & & \text{Hom}(\wedge^2 TM, TM) \end{array}$$

It is convenient to use the coframe θ to fix a trivialization of TM so that \hat{c} becomes a map

$$\hat{c} : X \rightarrow \text{Hom}(\wedge^2 \mathbb{R}^n, \mathbb{R}^n).$$

Finally, if we define n vector fields on X through

$$F_i = (\kappa_\theta)_* \frac{\partial}{\partial \theta_i} \in \mathfrak{X}(X),$$

then we can describe the classifying Lie algebroid $A_\theta \rightarrow X$ of θ explicitly as follows. We take A_θ to be the trivial vector bundle $A_\theta = X \times \mathbb{R}^n$ and define its structure by

$$\begin{aligned} [e_i, e_j] &= -\hat{c}(e_i \wedge e_j) \\ \sharp(e_i) &= F_i \end{aligned}$$

where $\{e_1, \dots, e_n\}$ denotes de canonical basis of sections.

REFERENCES

- [1] R. L. Bryant. Bochner-Kähler metrics. *J. of Amer. Math. Soc.*, 14(3):623–715, 2001.
- [2] A. Cannas da Silva and A. Weinstein. *Geometric models for noncommutative algebras*, volume 10 of *Berkeley Mathematics Lecture Notes*. American Mathematical Society, Providence, RI, 1999.
- [3] E. Cartan. Sur la structure des groupes infinis de transformations. *Ann. Éc. Norm.*, 3:153–206, 1904.
- [4] M. Crainic. Differentiable and algebroid cohomology, Van Est isomorphism, and characteristic classes. *Comment. Math. Helv.*, 78:681–721, 2003.
- [5] M. Crainic and R. L. Fernandes. On integrability of Lie brackets. *Annals of Mathematics*, 157:575–620, 2003.
- [6] M. Crainic and R. L. Fernandes. Secondary characteristic classes of Lie algebroids. In *Quantum field theory and noncommutative geometry*, volume 662 of *Lecture Notes in Phys.*, pages 157–176. Springer, Berlin, 2005.
- [7] M. Crainic and R. L. Fernandes. Lectures on integrability of Lie brackets. *Geometry & Topology Monographs*, 17:1–94, 2010.
- [8] S. Evans, J. H. Lu, and A. Weinstein. Transverse measures, the modular class and a cohomology pairing for Lie algebroids. *Quart. J. Math.*, 50(Ser. 2):417–436, 1999.
- [9] R. L. Fernandes. Lie algebroids, holonomy and characteristic classes. *Adv. Math.*, 170(1):119–179, 2002.
- [10] R. L. Fernandes and I. Struchiner. The classifying Lie algebroid of a geometric structure II: G -structures and examples. *preprint in preperation*.
- [11] S. Kobayashi. *Transformation Groups in Differential Geometry*. Springer-Verlag, 1972.
- [12] Y. Kosmann-Schwarzbach, C. Laurent-Gengoux, and A. Weinstein. Modular class of Lie algebroid morphisms. *Preprint available at arXiv:0712.3021 [math.DG]*, pages 1–33, 2008.
- [13] P. J. Olver. *Equivalence, Invariants and Symmetry*. Cambridge University Press, 1995.
- [14] P. J. Olver. Non-associative local Lie groups. *J. Lie Theory*, 6:23–51, 1996.
- [15] R. S. Palais. A global formulation of the Lie theory of transformation groups. *Mem. Amer. Math. Soc.*, 22:iii+123pp, 1957.
- [16] Proceedings of the "Rencontres Mathématiques de Glanon". *Symplectic Connections via Integration of Poisson Structures*, 2002.
- [17] R. W. Sharpe. *Differential Geometry: Cartan's Generalization of Klein's Erlangen Program*, volume Volume 166 of *Graduate Texts in Mathematics*. Springer, 1997.
- [18] I. M. Singer and S. Sternberg. The infinite groups of Lie and Cartan. *J. D'Anal. Math.*, 15:1–113, 1965.
- [19] S. Sternberg. *Lectures on Differential Geometry*. Prentice-Hall, 1964.
- [20] I. Struchiner. *The Classifying Lie Algebroid of a Geometric Structure*. PhD thesis, University of Campinas - UNICAMP, 2009.