

Geometrical Structures of Space-Time in General Relativity (Poster)*

I. SÁNCHEZ RODRÍGUEZ

September, 2007

Abstract

Space-Time in general relativity is a dynamical entity because it is subject to the Einstein field equations.

From the point of view of differential geometry, the space-time is a manifold with a Lorentzian metric. The space-time metric provides different geometrical structures: conformal, volume, projective . . .

A deep understanding of the geometrical structures has consequences on the dynamical role played by geometry. We explain these geometrical structures, establishing relationships among them and clarifying the meaning of associated geometric magnitudes.

Recently, some of my research [1,2] have been taken into consideration for one of the lines of thought about quantum gravity [3,4]. This poster is a set of my latest reflections and conclusions about the applications of my work to physics.

References

1. SÁNCHEZ-RODRÍGUEZ,I., “*Intersection of G-structures of first or second order*”. In O. Kowalski (ed.) et al., *Differential geometry and its applications. Proceedings of the 8th International Conference, Opava, Czech Republic, August 27-31, 2001*, 135-140, Opava: Silesian University at Opava, Math. Publ. Vol. 3, 2001.
2. SÁNCHEZ RODRÍGUEZ,I., “*On the intersection of geometrical structures*”. In M. A. Cañadas-Pinedo, (ed.) et al., *Lorentzian Geometry-Benalmádena 2001. Benalmádena (Málaga) Spain, November 14-16, 2001*, 239-246, Publicaciones de la Real Sociedad Matemática Española, Vol. 5, 2003.
3. STACHEL, JOHN, “*Structure, individuality and quantum gravity*”. In D. Rickles (ed.) et al., *The Structural Foundations of Quantum Gravity*, 53-82, Oxford: Oxford University Press, Clarendon, 2006.
4. STACHEL, J., “*Prolegomena to any future Quantum Gravity*”. ArXiv:gr-qc/0609108 v1, 23 Sep 2006.

* Poster presented at the XVI International Fall Workshop on Geometry and Physics, 5-8 September 2007, Lisbon (Portugal).

1 Introduction

The space-time of general relativity (GR) is a 4-dimensional manifold M , with a C^∞ atlas \mathcal{A} . The atlas is the *differential structure* of our space-time.

The *equivalence principle* of GR establishes the invariance by diffeomorphisms. This let us to think that a *physical event* is not a point, but a geometrical structure on a neighborhood.

The *fundamental geometrical structures* that we consider defined in the space-time are:

- **Volume** (4-form)
- **Conformal structure** (Lorentzian)
- **Metric** (Lorentzian)
- **Linear connection** (symmetric)
- **Projective structure**

They are defined in terms of the *most primitive* differential structure, via the concept of G -structure. Volume, conformal and metric are *first order G -structures*. But linear connection and projective are *second order G -structures*.

For certain G 's, *classified in* [8], every first order G -structure lead to a unique second order structure, named its *prolongation*. This is the case for the volume, metric and conformal structures.

2 Frame bundles

The *r -th order frame bundle* $\mathcal{F}^r M$ is a quotient space in a subset of \mathcal{A} ([12, p.38]). An *r -frame*, $j^r \varphi \in \mathcal{F}^r M$ is an r -jet at 0, where $x = \varphi^{-1}$ is a chart with 0 as a target.

The first order frame bundle $\mathcal{F}^1 M$ is usually identified with the *linear frame bundle* LM .

Let LLM be the linear bundle of LM . There is a *canonical inclusion* $\mathcal{F}^2 M \hookrightarrow LLM$, $j^2 \varphi \mapsto j^1 \tilde{\varphi}$, where $\tilde{\varphi}$ is the diffeomorphism induced by φ , between neighborhoods of $0 \in \mathbf{R}^{n+n^2}$ and $j^1 \varphi \in LM$ ([10], [12, p.50]).

Let $J^1 LM$ be the bundle of *1-jets of (local) sections* of LM and s be a section of LM . Each $j_p^1 s$ is characterized by the *transversal n -subspace* $H_l = s_*(T_p M) \subset T_l LM$.

Then, there is also a *canonical inclusion* $J^1LM \hookrightarrow LLM$, $j_p^1s \mapsto z$, where z is the basis of T_lLM , whose first n vectors span H_l and correspond to the usual basis of \mathbf{R}^n , via the *canonical form of LM* (see [6]).

By the previous canonical maps, it happens that \mathcal{F}^2M is mapped one to one into the subset of J^1LM , corresponding with the *torsion-free transversal n -subspaces* in TLM .

Theorem 1. *We have the canonical embeddings:*

$$\mathcal{F}^2M \hookrightarrow J^1LM \hookrightarrow LLM$$

3 Structural groups

Each \mathcal{F}^rM is a principal bundle respect to the group G_n^r of *r -jets at 0 of diffeomorphisms of \mathbf{R}^n* , $j_0^r\phi$, with $\phi(0) = 0$.

The group G_n^1 is identified with $GL(n, \mathbf{R})$. Then, there is a canonical inclusion of G_n^1 into G_n^r , if we take the r -jet at 0 of every linear map of \mathbf{R}^n . Furthermore, G_n^r is the *semidirect product* of G_n^1 with a nilpotent normal subgroup (see [13] for details).

Let's see *the structure for G_n^2* . We consider the underlying additive group of the vector space S_n^2 of symmetric bilinear maps of $\mathbf{R}^n \times \mathbf{R}^n$ into \mathbf{R}^n . There is a monomorphism $\iota: S_n^2 \rightarrow G_n^2$ defined by $\iota(s) = j_0^2\phi$ with $s = (s_{jk}^i)$ and $\phi(u^i) := (u^i + \frac{1}{2}s_{jk}^i u^j u^k)$.

Theorem 2. *We obtain the split exact sequence of groups:*

$$0 \rightarrow S_n^2 \xrightarrow{\iota} G_n^2 \xrightarrow{\cong} G_n^1 \rightarrow 1$$

It makes G_n^2 *isomorphic to the semidirect product* $G_n^1 \rtimes S_n^2$, whose multiplication rule is $(a, s)(b, t) := (ab, b^{-1}s(b, b) + t)$. The isomorphism is given by $j_0^2\phi \mapsto (D\phi|_0, D\phi|_0^{-1}D^2\phi|_0)$.

For a linear group G , let \mathfrak{g} denote the Lie algebra of G . The *first prolongation of \mathfrak{g}* is defined by $\mathfrak{g}_1 := S_n^2 \cap L(\mathbf{R}^n, \mathfrak{g})$. We obtain that $G \times \mathfrak{g}_1$ *is a subgroup* of $G_n^1 \rtimes S_n^2$, and hence, a subgroup of G_n^2 (see more details in [1]).

4 G -structures

We define an *r -th order G -structure* on M as a *reduction* of \mathcal{F}^rM to a subgroup $G \subset G_n^r$ ([10]). The idea of *geometrical structure* on M concerns the classification of charts in \mathcal{A} , *when choosing the meaningful classes* guided by an structural group.

We exemplify the concept of a G -structure studying a volume on a manifold, which rarely is treated this way (see [3]).

Let's define *volume on M* as a first order G -structure V , with $G = \text{SL}_n^\pm := \{a \in \text{GL}(n, \mathbf{R}): |\det(a)| = 1\}$. For an orientable M , V has two components for two $\text{SL}(n, \mathbf{R})$ -structures, for two equal, except sign, *volume n -forms*. For a general M , volume corresponds to *odd type n -form*, as in [4, pp. 21-27].

From *principal bundle theory* ([9]), SL_n^\pm -structures are the sections of the *bundle associated* with LM and the left action of G_n^1 on $\text{G}_n^1/\text{SL}_n^\pm$. This is the *volume bundle*, \mathcal{VM} . Furthermore, the sections of \mathcal{VM} correspond to *G_n^1 -equivariant functions* f of LM to $\text{G}_n^1/\text{SL}_n^\pm$. The equivariance condition is $f(la) = |\det a|^{-\frac{1}{n}} I_n \cdot f(l)$, $\forall a \in \text{G}_n^1$. We have bijections:

$$\text{Volumes on } M \longleftrightarrow \text{Sec } \mathcal{VM} \longleftrightarrow C_{\text{eq}}^\infty(LM, \text{G}_n^1/\text{SL}_n^\pm)$$

The isomorphisms $\text{G}_n^1/\text{SL}_n^\pm \simeq \text{H}_n$, with $\text{H}_n := \{kI_n: k > 0\}$ and $\text{H}_n \simeq \mathbf{R}^+$, the multiplicative group of positive numbers, allow to represent a volume as an *(odd) scalar density* on M .

5 Second order structures

We can view a *symmetric linear connection (SLC) on M* as a G_n^1 -structure of second order. An SLC is also the image of an *injective homomorphism* of LM to \mathcal{F}^2M ([10]).

From the *principal bundle theory*, SLC's on M are sections of the *SLC bundle*, \mathcal{DM} , associated with \mathcal{F}^2M and the action of G_n^2 on $\text{G}_n^2/\text{G}_n^1 \simeq \text{S}_n^2$. Furthermore, each SLC, ∇ , corresponds to a *G_n^2 -equivariant function* $f^\nabla: \mathcal{F}^2M \rightarrow \text{S}_n^2$, verifying $f^\nabla(z(a, s)) = a^{-1}f^\nabla(z)(a, a) + s$. We have the bijections:

$$\text{SLC's on } M \longleftrightarrow \text{Sec } \mathcal{DM} \longleftrightarrow C_{\text{eq}}^\infty(\mathcal{F}^2M, \text{S}_n^2)$$

Given two SLC's, ∇ and $\widehat{\nabla}$, *the difference* $f^\nabla - f^{\widehat{\nabla}}$ verifies $z(a, s) \mapsto a^{-1}(f^\nabla(z) - f^{\widehat{\nabla}}(z))(a, a)$. Then, it is projectable to a function $f: LM \rightarrow \text{S}_n^2$ verifying $f(la) = a^{-1}f(l)(a, a)$, which corresponds to a *symmetric $\binom{1}{2}$ -tensor* $\rho = (\rho_{jk}^i)$ on M .

A *projective structure (PS)* is a set of SLC's which have the same family of *pregeodesics*. This is the cornerstone to understand the *freely falling bodies* in GR ([5]).

We define a PS on M as a *second order $\text{G}_n^1 \times \mathfrak{p}$ -structure*, \mathbf{Q} , with $\mathfrak{p} := \{s \in \text{S}_n^2: s_{jk}^i = \delta_j^i \mu_k + \mu_j \delta_k^i, \mu = (\mu_i) \in \mathbf{R}^{n*}\}$.

Now, for two SLC *included* in the same PS, i.e. $\nabla, \widehat{\nabla} \subset Q$, the tensor ρ , expressing *their difference*, is determined by the contraction $C(\rho) = (\rho_{si}^s)$, which is *an 1-form* on M .

6 Prolongations

Let B a *first order G -structure*. A connection in B is a distribution H of transversal n -subspaces, $H_l \subset T_l B$. If the subspaces are *free-torsion*, these determine a *second order G -structure*, whose G_n^1 -*extension* ([7, p.206]) is a SLC on M . Then, we say that B *admits an SLC*. Let us give two examples:

- An SLC and a parallel volume is an *equiaffine structure* on M ([11]); hence, it is a second order SL_n^\pm -structure.
- An SLC compatible with a conformal structure is a *Weyl structure*; hence, it is a second order $CO(n)$ -structure ([2]).

Theorem 3. *Let $B \subset LM$ a G -structure, admitting an SLC. Then, the set of 2-frames, corresponding with torsion-free transversal n -subspaces which are included in TB , is a *reduction of $\mathcal{F}^2 M$ to $G \times \mathfrak{g}_1$* . It is named the *prolongation of B* and denoted by B^2 . (For a proof, see [12, p.150-155]).*

Let us give a well known example: if B is an $O(n)$ -structure, B^2 is isomorphic to B on account of $\mathfrak{o}(n)_1 = \{0\}$; this explain the uniqueness of Levi-Civita connection.

There is an important theorem ([8]) *classifying the groups G* such that *every G -structure admits an SLC*: only the groups of *volume, metric and conformal* structures, and a class of groups preserving an 1-dimensional distribution, have this property.

7 Concluding remarks

We have done a unified description of the geometrical structures that have been used by GR to define intrinsic properties of the space-time. The unifying criterion we used for it, *not only is natural in the sense that geometric objects are sections of bundles associated with the $\mathcal{F}^r M$ frame bundles* ([13]), *but that the objects themselves are reductions of $\mathcal{F}^r M$* . We have not considered a *linear connection with torsion* because it is a section of an associated bundle of $\mathcal{F}^2 M$, but not a reduction.

We have tried to clarify the relationships between the structures involved. Only simple relations, such as intersection, inclusion, reduction and

extension, have been used for it, on account of the previous *prolongation* of G -structures admitting SLC. For instance, it follows readily from the last section that the classical *equiaffine* or *Weyl structures* can be defined as the intersection of an SLC with the prolongation of a volume or a conformal structure, respectively.

Acknowledgments

The author would like to thank the IFWGP'07 organizers for their kind invitation to attend this Workshop and to present this poster. I wishes to acknowledge gratefully Dr. John Stachel the attention paid to my work. This work has been partially supported by the Junta de Andalucía P.A.I.: FQM-324.

References

- [1] E. AGUIRRE DABÁN, I. SÁNCHEZ RODRÍGUEZ, “*Explicit formulas for the 3-jet lift of a matrix group. Applications to conformal geometry*”. In: A. Pereira (ed.) et al., *Proc.1st Int. Meeting on Geometry and Topology*, 191-205, Braga: Universidade do Minho, 1998. Available online: <http://www.emis.ams.org/proceedings/Braga97/>
- [2] R.A COLEMAN, H. KORTE, “*Spacetime G structures and their prolongations*”, *J. Math. Phys.* 22 (1981), 2598-2611.
- [3] M. CRAMPIN, D.J. SAUNDERS, “*Projective connections*”, *Journal of Geometry and Physics*, 57 (2007), 691727.
- [4] G. DE RHAM, *Variétés différentiables*, 2nd ed., Paris: Hermann, 1960.
- [5] J. EHLERS, F.A.E. PIRANI AND A. SCHILD, “*The geometry of free fall and light propagation*”. In: L. O’Raifeartaigh (ed.), *General Relativity*, 63-84, Oxford: Clarendon, 1972.
- [6] P.L. GARCIA, “*Connections and 1-jet fiber bundles*”, *Rend. Sem. Mat. Padova*, 47 (1972), 227-242.
- [7] W. GREUB, S. HALPERIN, R. VANSTONE, *Connections, Curvature and Cohomology (vol. II)*, New York: Academic Press, 1973.
- [8] S. KOBAYASHI, T. NAGANO, “*On a fundamental theorem of Weyl-Cartan on G -structures*”, *J. Math. Soc. Japan*, 17 (1965), 84-101.

- [9] S. KOBAYASHI, K. NOMIZU, *Foundations of Differential Geometry (vol. I)*, New York: John Wiley - Interscience, 1963.
- [10] S. KOBAYASHI, *Transformation Groups in Differential Geometry*, Heidelberg: Springer, 1972.
- [11] K. NOMIZU, T. SASAKI, *Affine differential geometry*, Cambridge University Press, 1994.
- [12] IGNACIO SÁNCHEZ RODRÍGUEZ, *Conexiones en el fibrado de referencias de segundo orden. Conexiones conformes*, Doctoral Thesis, Complutense University of Madrid, Madrid, 1994. Available online: <http://www.ugr.es/local/ignacios/>
- [13] C. L. TERNG, “Natural vector bundles and natural differential operators”, *Am. J. Math.* 100 (1978), 775-828.

Author address: (Ignacio Sánchez Rodríguez)

DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA,
FACULTAD DE CIENCIAS, UNIVERSIDAD DE GRANADA,
AVDA. FUENTENUEVA S/N, 18071-GRANADA, SPAIN
E-mail: ignacios@ugr.es
URL: <http://www.ugr.es/local/ignacios>